

# **Attention problems in relation to gestational age at birth and smallness for gestational age**

**Authors:** Suna Eryigit-Madzwamuse, PhD; Brighton, UK; Centre for Health Research, University of Brighton.

Dieter Wolke, PhD; Coventry, UK; Department of Psychology, University of Warwick; Warwick Medical School, University of Warwick

**Paper presentation information:** An earlier version of this paper was presented at the 54<sup>th</sup> annual meeting of the European Society for Paediatric Research, Porto, Portugal, October 10-13, 2013.

**Corresponding author:** Dieter Wolke, Department of Psychology, University of Warwick, Coventry CV4 7AL, United Kingdom, work phone: +44 2476573217, home/cell phone: +44 7824358737 fax: +44 2476524225, [D.Wolke@warwick.ac.uk](mailto:D.Wolke@warwick.ac.uk)

**Short title:** Attention problems and gestational age

**Abstract- word count:** 243

**Main text- word count:** 2902

Eryigit-Madzwamuse, S., & Wolke, D. (2015). Attention problems in relation to gestational age at birth and smallness for gestational age. *Early Human Development*, 91(2), 131-138. doi: <http://dx.doi.org/10.1016/j.earlhumdev.2015.01.004>

## Abstract

**Background:** While it is well established that very preterm birth (gestational age at birth < 32 weeks) is related to increased attention problems, there is still considerable uncertainty of the effects of moderate or late preterm birth or smallness for gestational age (SGA) on attention regulation.

**Aims:** To investigate the impact of gestational age at birth and SGA, birth on child attention problems.

**Study design:** Prospective longitudinal cohort study.

**Subjects:** A total of 1435 children sampled from the Bavarian Longitudinal Study (BLS).

**Outcome measures:** Main outcome variables were parent-reported attention problems and examiner-reported attention skills at 6 and 8 years. Predictors were linked to attention outcomes using hierarchical regression analyses.

**Results:** Gestational age at birth ranged from 25 weeks to 41 weeks. We found a quadratic effect of gestational age on attention problems ( $\beta_{6\text{years}}=0.161$ , 95% CI= 0.085; 0.236;  $\beta_{8\text{years}}=0.211$ , 95% CI= 0.135; 0.287), and attention skills at 6 and 8 years ( $\beta_{6\text{years}}=-0.178$ , 95% CI= -0.252; -0.104;  $\beta_{8\text{years}}=-0.169$ , 95% CI= -0.243; -0.094). Elective caesarean birth did not predict child attention. In adjusted models, SGA was an additional risk factor for attention problems ( $\beta=0.080$ , 95% CI= 0.026; 0.134), and attention skills ( $\beta=-0.091$ , 95% CI=-0.143; -0.039) at 6 years but not at 8 years after adjusting for child sex and family SES.

**Conclusion:** Adverse effects on attention are disproportionately higher at early gestations. In contrast, the impact of SGA status was found to be similar at all gestational ages but disappeared by 8 years.

**Keywords:** child attention, preterm, gestational age at birth, elective caesarean birth, small for gestational age

## 1. Introduction

Attention skills are a fundamental prerequisite of learning and attention problems predict lower academic success (1, 2). Attention skills have been regularly reported to be impaired in children and adolescents who were born very preterm (gestational age at birth <32 weeks) (3-6). In contrast, findings are inconsistent for children born moderately or late preterm (gestational age at birth between 32 and 36 weeks) (3, 7-10).

Up to a third of preterm births are iatrogenic because of concerns about maternal or fetal health (11). Elective caesarean birth is common in iatrogenic births, and this mode of delivery has been associated with adverse outcomes in the neonatal period (12-15) and beyond (8, 16). Talge and her colleagues (8) reported a significant association between elective caesarean birth and attention problems in children born late preterm (34- 36 weeks).

Fetal growth restriction (FGR), commonly associated with a variety of pregnancy complications (i.e., preeclampsia), is a common reason for elective caesarean birth (8). Smallness for gestational age (SGA: i.e. a birth weight below 10<sup>th</sup> percentile) has been used as a proxy for FGR (17). SGA has been reported to significantly predict attention related problems in some studies (6, 10, 18) but not in others (8, 19, 20). Thus, while the link between very preterm birth and childhood attention problems is well established, the impacts of moderate/late preterm birth, SGA and elective caesarean birth are still uncertain.

The objective of this study was to investigate the impact of gestational age at birth, and SGA on child attention at 6 and 8 years across the full gestation spectrum. Firstly, we examined the impact of gestational age on child attention. We hypothesized that gestational age at birth is associated with child attention at school age (6 and 8 years) independent of whether reported by parents or examiner. We also hypothesized that the relation of attention with gestational age at birth would be non-linear because attention difficulties are disproportionately higher after very preterm birth than previously reported after later preterm birth. Secondly, we examined whether SGA additionally predicts attention outcomes while controlling for potential confounders such as

prenatal complications, elective caesarean birth, head circumference in infancy and family socio-economic status (hereafter family SES).

## 2. Materials and Methods

### 2.1. Study Sample

Details of the design of the Bavarian Longitudinal Study have been described previously (21). Ethical approval of this study was granted by the Ethical Review Board of the Rheinische Friedrich Wilhelms University, Medical Faculty on 18 August 2009 (reference # 159/09). Informed consent was provided by the parents within 48 hours of their child's birth. Subjects were children born alive in a geographically defined area of Southern Bavaria (Germany) between February 1985 and March 1986 and who required admission to children's hospitals within the first 10 days after birth (N=7505; 10.6% of all live births). Healthy infants who were born in the same obstetric hospitals, cared for on normal postnatal wards were recruited as controls (N=916), resulting in a total study sample of 8421 newborns. Of the 8421 participants, 1513 children were selected for follow-up at 6 and 8 years of age according to the following criteria: 1. either very preterm (<32 weeks gestation) or very low birth weight (<1500g birth weight); 2. a subsample of children born at  $\geq 32$  weeks gestation randomly selected within the stratification factors gender, family SES (low, medium and high) and degree of neonatal risk (none, low, moderate, high and very high) (22).

Of the eligible children, 1475 participated in the follow-up at both 6 and 8 years. Children who did not attend the assessments either at 6 years or at 8 years (N=38) did not differ from children who attended assessments at both time points (N=1475) in terms of gestational age, prenatal complications, mode of delivery, SGA status and head circumference but were more often of low family SES ( $\chi^2(1) = 21.315, p < 0.001$ ). Furthermore, we excluded an additional 40 children who were born post-term (42 or more completed weeks of gestation) because post term birth has different causes and is associated with an increased risk of developmental abnormalities (23). Table 1 shows the characteristics of the final sample with gestational age at birth ranging from 25 weeks to 41 weeks (N=1435).

### 2.2. Measures

Gestational age was determined from a combination of maternal reports of the last menstrual period and serial ultrasound fetal biometry during pregnancy. Prenatal data were coded from the standard Bavarian perinatal survey (24) assessed prospectively and included 14 items (e.g. preeclampsia) (Table 1). Higher scores on this scale indicate more complications with a range from 0 (best condition) to 14 (most risky condition) (25).

Mode of delivery was coded in the Bavarian perinatal survey (24) as elective caesarean birth, emergency caesarean birth or vaginal birth. For the purposes of the current study we reclassified mode of delivery as elective caesarean birth versus delivery following labour (vaginal birth or emergency caesarean birth). Children with birth weight less than the gender specific 10<sup>th</sup> percentile for gestational age according to a perinatal survey of all Bavarian newborns (24) were classified as small for gestational age (SGA); 376 children were SGA and 1059 children were born with a birth weight appropriate for their gestational age (AGA). Figure 1 shows the frequency of children who were born small across the whole gestation spectrum. SGA was most frequent between 32 weeks and 37 weeks with a range from 35% to 50% of total births at each week.

The primary outcome of this study was the level of attention of the children reported by parents and examiners separately at the ages of 6 and 8 years. Mothers rated their children's attention problems using the attention problems scale of the Child Behaviour Check List (CBCL(26)). Higher scores on this 11 item scale indicate more attention problems.

Examiners reported on children's attention skills in two ways. First, the assistant psychologist rated children's behaviour during a cognitive assessment that lasted on average 80 minutes using the Tester Ratings of Child Behaviour (TRCB (1, 27)). Six of 13 items were combined to create a Task Orientation subscale including attention, robustness/endurance, demandingness (reversed score), cooperativeness, compliance and difficulty (reversed score). Inter-class correlations in inter-rater reliabilities ranged from 0.63 to 0.97. Second, adapting a consensus model, the whole research team (psychologist, assistant psychologist and paediatrician) evaluated children's behaviour based on their observations during the whole assessment day. They rated children's behaviours using three items of the TRCB: attention, robustness/endurance and demandingness (reversed score) (1). An index score of attention

was computed using the Team ratings. Strong correlations were found between the TRCB and Team ratings ( $r= 0.79$ , and  $0.78$  at 6 and 8 years, respectively;  $p<0.001$ ). Thus, we computed a mean score in order to reflect examiner-reported attention skills. Higher scores on this scale from 1 (very low) to 9 (very high) indicate better attention skills.

*Potential confounders.* Both head circumference (6, 28) and family SES (20, 29) have been previously reported to predict attention levels in children. Head circumference is a proxy measure of brain growth and was assessed at 5 months (corrected for gestation at birth) by research nurses who measured head circumference twice and the mean of both measures was recorded. Family SES was a weighted composite score of parents' education and occupation and grouped as high, middle and low according to predefined cut-off points .

### 2.3. Statistical Analysis

We used SPSS 21.0 to analyse the data. First, we examined all the variables for outliers and normality. Next, we compared all included variables between AGA and SGA groups using One-Way ANOVA for continuous variables and  $\chi^2$  test for categorical variables. Prior to testing the first research question about the effect of gestational age, we fitted curve estimations and found the quadratic model to be the best fitting model. Next, we fitted regression analyses with gestational age and its quadratic term as predictors for each outcome variable separately. Prior to regression, we centred gestational age at 37 weeks as an attempt to decrease co-linearity between the gestational age linear term and the gestational age quadratic term. To illustrate the quadratic relation, we fitted regression lines from week 25 to week 41 with their 95% confidence intervals.

In order to address the second research question regarding the effects of elective caesarean birth and SGA, we fitted hierarchical regression models where we entered prenatal complications and elective caesarean birth into a third model after gestational age and its quadratic term; SGA in the fourth model; and head circumference in the fifth model. In the final model, we adjusted the analyses according to family SES.

## 3. Results

Table 1 reports the sample characteristics. On average there was a significant decrease in parent-reported attention problems from 6 years to 8 years ( $t_{1335}=18.973$ ,  $p<0.001$ ) and there

was a significant increase in examiner-reported attention skills ( $t_{1310}=23.759$ ,  $p<0.001$ ) from 6 years to 8 years.

Table 1 also shows the comparisons across the two birth weight groups, AGA and SGA. AGA children had, on average, higher gestational age at birth, fewer prenatal complications, were born more frequently following labour and had higher head circumferences at 5 months compared to the SGA group. Furthermore, SGA children were more often of low SES families.

### 3.1. Is the relation between gestation and child attention linear or non-linear?

We examined the relationship between gestational age and attention outcomes at 6 and 8 years. Model 1 and Model 2 in Table 2 show the standardized beta coefficients and their 95% confidence intervals for parent-reported attention problems. The gestational age quadratic term was a significant predictor at both 6 and 8 years. As shown in Figure 2a and 2b, the relation between gestational age and parent-reported attention problems is concave, indicating higher attention problems with every week of gestation either before or after 37 weeks. However, at gestational ages later than 37 weeks, differences in attention problems per week of gestation were very small.

The effect of gestational age at birth on attention was similar on the examiner-reported attention skills (see Model 1 and Model 2 in Table 3). Both the linear and quadratic terms were significant predictors at both 6 years and 8 years. As can be seen in Figure 2c and 2d, the relation between gestational age at birth and examiner-reported attention skills is also concave but downwards, indicating lower attention skills both before and after 37 weeks but very small differences per week of gestation after 37 weeks indicating no significant difference between 37 and 41 weeks.

### 3.2. What is the effect of SGA on attention in childhood?

We fitted hierarchical regression models separately for each outcome variable (see Table 2 & Table 3). Adjusting for gestational age, prenatal complications, and elective caesarean birth being SGA was consistently a risk factor for attention problems or reduced attention skills. Further, we examined the interaction between gestational age and SGA status and found no interaction, indicating that the impact of being SGA was similar across the whole gestation spectrum. The impact of being SGA was attenuated but was still significant once

adjusted for head circumference at 5 months.. However, its impact at 8 years disappeared when adjusted for child sex and family SES.

#### **4. Discussion**

We found a significant non-linear relationship between gestational age at birth and attention outcomes in childhood regardless of the source of data (parents, examiners who were blind to preterm status) or age at assessment (6 and 8 years). The non-linear relationship remained significant after adjusting for prenatal complications, whether born after labour or by elective caesarean, SGA status, head circumference and family SES. Furthermore, being SGA was an additional risk factor for poorer attention in childhood across the full spectrum of gestation; however its impact disappeared by 8 years.

The strengths of this study are its prospective design and large sample; inclusion of the full spectrum of gestational age, with a focus on gestational age in weeks rather than aggregated periods and using multiple sources for child outcomes at different ages. There are also limitations. Induction of labour is a factor that has been associated with improved maternal and/or fetal condition at birth (30, 31). We were unable to examine the impact of induction as induction and acceleration after labour onset were coded as one variable. Second, SGA was used as a proxy of FGR. However, some newborns are constitutionally small rather than growth restricted and their prognosis is less likely to be influenced by SGA status (32). We were unable to differentiate constitutionally small newborns from FGR newborns due to lack of data on pre-pregnancy maternal weight which is a crucial factor when defining constitutional smallness (33). Third, head circumference was employed as a proxy measure of brain volume. Previous research has indicated that head circumference accurately predicts brain volume (34) with a correlation with brain volume in 1 to 6 year old children in MRI of  $r = 0.93$  (35). Lastly, due to improvements in neonatal care and increased survival of extremely preterm children the rates of attention problems or reduced attention skills may be similar or may even have increased in recent cohorts of preterm children (3). Research in recent cohorts is needed in order to examine how changes in care for larger and smaller preterm children may have altered outcome in attention regulation. Previous studies of attention problems were limited to specific periods of gestation, either the very preterm period, (3) late preterm period (7) or term-born

children only (36). Very few analysed the effect on attention deficit/hyperactivity disorder (ADHD) across the whole gestation spectrum and reported inconsistent findings (10, 20). Moreover, these studies did not distinguish between inattention (ADD) problems and hyperactivity/impulsivity problems (ADHD); there is substantial empirical evidence for the risk of inattention, rather than hyperactivity symptoms of ADHD in relation to prematurity (1). We analysed the whole gestation spectrum and found a significant non-linear association between gestational age at birth and child attention. To illustrate the quadratic effect at age 6 years, the decrease in attention problems with one week increase in gestation at 31 weeks (i.e., the difference between 31 weeks and 32 weeks) was 8.6 times higher than the decrease in attention problems with one week increase in gestation at 37 weeks (i.e., the difference between 37 and 38 weeks). Similarly, the increase in attention skills for one week increase in gestation at 31 weeks was 3.5 times higher than the increase in attention skills for one week increase in gestation at 37 weeks. Our study provides strong support for an increased risk of inattention with decreasing gestation at birth. This relationship was found to be non-linear with increasing adverse impact on attention with each week of gestation the lower the gestation.

Consistent with previous studies (1, 3-6), we found that very preterm children still have the lowest attention skills at 8 years of age, showing the importance of reducing the rate of preterm birth. Unfortunately, to date, no interventions designed to prolong pregnancy have been shown to result in an improved long term outcome. This may be because prolonging pregnancy without ameliorating the adverse intrauterine environment associated with impending preterm birth simply keeps the baby longer in a damaging environment, thus negating any improvement consequent upon being born at a later gestational age. This emphasizes the importance of addressing the factors leading to preterm birth (e.g. infection, malnutrition, social disadvantage, stress, drugs, induced multiple pregnancy from IVF, hypertension and diabetes) rather than trying to prolong the duration of an abnormal pregnancy (37).

In unadjusted crude analysis, we found a significant effect of elective caesarean birth on attention outcomes (data not shown) but this effect became non-significant once adjusted for gestational age. Previously, adverse outcomes of elective caesarean during childhood (8) have been reported when gestational age was not taken into account. In our sample, the average

gestational age at birth in the elective caesarean group was 4 weeks earlier than the gestational age at birth in the labouring group. Therefore, the significant effect of elective caesarean birth found in previous studies might be partly explained by differences in gestational ages at birth.

The current SGA results are consistent with previous findings suggesting that SGA is an additional risk factor for attention problems or reduced attention skills (6, 10, 38). Our results shed light on the inconsistent findings regarding the impact of SGA across the full gestation spectrum (10, 39); SGA as defined in our study has similar adverse impact across the whole gestation range. The impact of SGA status is also only partially linked to brain size. After adjusting for head circumference as an indicator of brain size, the impact of SGA on child attention was attenuated but was still significant, suggesting that SGA might be linked to both reduction in brain volume and alteration in brain structure (39). Once child sex and family SES were controlled the impact of SGA on 8 years attention problems and reduced attention skills disappeared; however, no interaction between SGA and child sex or between SGA and family SES were found (data not shown).

To conclude, our study suggests that attention problems are not limited to children born very preterm but gestational age at birth matters for all children below 37 weeks of gestation. Fetal growth restriction is amongst the most common reasons for elective caesarean birth and one reason for shortened gestation. Future research should concentrate on ways of improving fetal development and preventing early deliveries by improving maternal health; this will in turn improve neonatal and childhood development.

**Acknowledgement:** We are grateful to funders who supported this study, all research assistants, psychologists and paediatricians who collected the data and to the parents and their children for their participation. We are grateful to Professor P J Steer for helpful suggestions while the manuscript was being written, and for assistance with editing the text.

**Conflict of interest:** The authors report no conflict of interest

**Role of the Funding Source:** This study was funded by grants PKE24, JUG14, 01EP9504 and 01ER0801 from the German Federal Ministry of Education and Science (BMBF). The contents are solely the responsibility of the authors and do not necessarily represent the official view of BMBF. Information on BMBF is available on <http://www.bmbf.de/en/>.

### **Highlights**

- We model attention problems and skills across the whole gestation spectrum.
- We compare both parent and examiner reports of attention and at 6 and 8 years.
- We find a non-linear relation between gestational age and attention.
- The impact of being born SGA is similar across the whole gestation spectrum.

## References

1. Jaekel J, Wolke D, Bartmann P. Poor attention rather than hyperactivity/impulsivity predicts academic achievement in very preterm and full-term adolescents. *Psychol Med*. 2013;43(1):183-96.
2. Duncan GJ, Claessens A, Huston AC, Pagani LS, Engel M, Sexton H, et al. School readiness and later achievement. *Dev Psychol*. 2007;43(6):1428-46.
3. Johnson S, Wolke D. Behavioural outcomes and psychopathology during adolescence. *Early Human Development*. 2013;89(4):199-207.
4. Aarnoudse-Moens CSH, Weisglas-Kuperus N, van Goudoever JB, Oosterlaan J. Meta-Analysis of Neurobehavioral Outcomes in Very Preterm and/or Very Low Birth Weight Children. *Pediatrics*. 2009;124(2):717-28.
5. Bhutta AT, Cleves MA, Casey PH, Cradock MM, Anand KJS. Cognitive and behavioral outcomes of school-aged children who were born preterm - A meta-analysis. *Jama-Journal of the American Medical Association*. 2002;288(6):728-37.
6. Hall J, Jaekel J, Wolke D. Gender distinctive impacts of prematurity and small for gestational age (SGA) on age-6 attention problems. *Child Adolesc Ment Health*. 2012;17(4):238-45.
7. de Jong M, Verhoeven M, van Baar AL. School outcome, cognitive functioning, and behaviour problems in moderate and late preterm children and adults: A review. *Seminars in Fetal & Neonatal Medicine*. 2012;17(3):163-9.
8. Talge NM, Holzman C, Van Egeren LA, Symonds LL, Scheid JM, Senagore PK, et al. Late-Preterm Birth by Delivery Circumstance and Its Association With Parent-Reported Attention Problems in Childhood. *J Dev Behav Pediatr*. 2012;33(5):405-15.
9. van Baar AL, Vermaas J, Knots E, de Kleine MJK, Soons P. Functioning at School Age of Moderately Preterm Children Born at 32 to 36 Weeks' Gestational Age. *Pediatrics*. 2009;124(1):251-7.

10. Heinonen K, Raikkonen K, Pesonen AK, Andersson S, Kajantie E, Eriksson JG, et al. Behavioural symptoms of attention deficit/hyperactivity disorder in preterm and term children born small and appropriate for gestational age: A longitudinal study. *BMC Pediatr.* 2010;10.
11. Saigal S, Doyle LW. Preterm birth 3 - An overview of mortality and sequelae of preterm birth from infancy to adulthood. *Lancet.* 2008;371(9608):261-9.
12. Chiossi G, Lai Y, Landon MB, Spong CY, Rouse DJ, Varner MW, et al. Timing of Delivery and Adverse Outcomes in Term Singleton Repeat Cesarean Deliveries. *Obstetrics & Gynecology.* 2013;121(3):561-9 10.1097/AOG.0b013e3182822193.
13. Laughon SK, Reddy UM, Sun L, Zhang J. Precursors for Late Preterm Birth in Singleton Gestations. *Obstetrics and Gynecology.* 2010;116(5):1047-55.
14. Kamath BD, Todd JK, Glazner JE, Lezotte D, Lynch AM. Neonatal Outcomes After Elective Cesarean Delivery. *Obstetrics and Gynecology.* 2009;113(6):1231-8.
15. Werner EF, Savitz DA, Janevic TM, Ehsanipoor RM, Thung SF, Funai EF, et al. Mode of Delivery and Neonatal Outcomes in Preterm, Small-for-Gestational-Age Newborns (vol 68, pg 9, 2013). *Obstetrical & Gynecological Survey.* 2013;68(2):151-3.
16. Walker D-M, Marlow N, Upstone L, Gross H, Hornbuckle J, Vail A, et al. The Growth Restriction Intervention Trial: long-term outcomes in a randomized trial of timing of delivery in fetal growth restriction. *American journal of obstetrics and gynecology.* 2011;204(1):34.e1-9.
17. Wollmann HA. Children Born Small for Gestational Age: Definitions and Etiology. In: Kiess W, Chernausek SD, editors. *Pediatric and Adolescent Medicine. Pediatric and Adolescent Medicine.* 132009. p. 1-10.
18. Gutbrod T, Wolke D, Soehne B, Ohrt B, Riegel K. Effects of gestation and birth weight on the growth and development of very low birthweight small for gestational age infants: a matched group comparison. *Archives of Disease in Childhood.* 2000;82(3):F208-F14.
19. Gustafsson P, KÄLIÉN K. Perinatal, maternal, and fetal characteristics of children diagnosed with attention-deficit–hyperactivity disorder: results from a population-based study

- utilizing the Swedish Medical Birth Register. *Developmental Medicine & Child Neurology*. 2011;53(3):263-8.
20. Lindstrom K, Lindblad F, Hjern A. Preterm Birth and Attention-Deficit/Hyperactivity Disorder in Schoolchildren. *Pediatrics*. 2011;127(5):858-65.
  21. Wolke D, Schmid G, Schreier A, Meyer R. Crying and Feeding Problems in Infancy and Cognitive Outcome in Preschool Children Born at Risk: A Prospective Population Study. *J Dev Behav Pediatr*. 2009;30(3):226-38.
  22. Jaekel J, Baumann N, Wolke D. Effects of Gestational Age at Birth on Cognitive Performance: A Function of Cognitive Workload Demands. *Plos One*. 2013;8(5).
  23. Boyle EM, Poulsen G, Field DJ, Kurinczuk JJ, Wolke D, Alfirevic Z, et al. Effects of gestational age at birth on health outcomes at 3 and 5 years of age: population based cohort study. *Br Med J*. 2012;344.
  24. Zander J, Holzmann K, Selbmann HK. [Data from the Bavarian perinatal survey on the problem of the incidence of cesarean section]. *Geburtshilfe und Frauenheilkunde*. 1989;49(4):328-36.
  25. Molfese VJ, Thomson B. OPTIMALITY VERSUS COMPLICATIONS - ASSESSING PREDICTIVE VALUES OF PERINATAL SCALES. *Child Development*. 1985;56(4):810-23.
  26. TM A. *The Manual for the Child Behavior Checklist 4- 18, 1991 Profile*. Burlington VT: University of Vermont, Department of Psychiatry; 1991.
  27. Wolke D, Skuse D, Mathisen B. BEHAVIORAL STYLE IN FAILURE-TO-THRIVE INFANTS - A PRELIMINARY COMMUNICATION. *Journal of Pediatric Psychology*. 1990;15(2):237-54.
  28. Cheong JLY, Hunt RW, Anderson PJ, Howard K, Thompson DK, Wang HX, et al. Head growth in preterm infants: Correlation with magnetic resonance imaging and neurodevelopmental outcome. *Pediatrics*. 2008;121(6):E1534-E40.

29. Ekeus C, Lindstrom K, Lindblad F, Rasmussen F, Hjern A. Preterm Birth, Social Disadvantage, and Cognitive Competence in Swedish 18- to 19-Year-Old Men. *Pediatrics*. 2010;125(1):E67-E73.
30. Koopmans CM, Bijlenga D, Groen H, Vijgen SMC, Aarnoudse JG, Bekedam DJ, et al. Induction of labour versus expectant monitoring for gestational hypertension or mild pre-eclampsia after 36 weeks' gestation (HYPITAT): a multicentre, open-label randomised controlled trial. *Lancet*. 2009;374(9694):979-88.
31. Stock SJ, Ferguson E, Duffy A, Ford I, Chalmers J, Norman JE. Outcomes of elective induction of labour compared with expectant management: population based study. *Br Med J*. 2012;344.
32. Resnik R. Intrauterine growth restriction. *Obstetrics and Gynecology*. 2002;99(3):490-6.
33. Gardosi J, Clausson B, Francis A. The value of customised centiles in assessing perinatal mortality risk associated with parity and maternal size. *Bjog-an International Journal of Obstetrics and Gynaecology*. 2009;116(10):1356-63.
34. Winick M, Rosso P. HEAD CIRCUMFERENCE AND CELLULAR GROWTH OF THE BRAIN IN NORMAL AND MARASMIC CHILDREN. *Journal of Pediatrics*. 1969;74(5):774-8.
35. Bartholomeusz HH, Courchesne E, Karns CM. Relationship between head circumference and brain volume in healthy normal toddlers, children, and adults. *Neuropediatrics*. 2002;33(5):239-41.
36. Yang S, Platt RW, Kramer MS. Variation in Child Cognitive Ability by Week of Gestation Among Healthy Term Births. *American Journal of Epidemiology*. 2010;171(4):399-406.
37. Steer PJ. The epidemiology of preterm labour--why have advances not equated to reduced incidence? *BJOG : an international journal of obstetrics and gynaecology*. 2006;113 Suppl 3:1-3.

38. Yang S, Fombonne E, Kramer MS. Duration of gestation, size at birth and later childhood behaviour. *Paediatric and Perinatal Epidemiology*. 2011;25(4):377-87.
39. Cornforth CM, Thompson JMD, Robinson E, Waldie KE, Pryor JE, Clark P, et al. Children born small for gestational age are not at special risk for preschool emotion and behaviour problems. *Early human development*. 2012;88(7):479-85.

Figure 1. Frequency of infants born with a birth weight small for their gestational age (SGA) and appropriate for gestational age (AGA) for each gestational ages in weeks.

Legend:

SGA status

 Appropriate for gestational age (AGA)

 Small for gestational age (SGA)

Figure 2. Predicted change in attention outcomes in relation to gestational age at birth (solid line with 95% confidence intervals in dotted line)

2a. Parent-reported attention problems at 6 years

2b. Parent-reported attention problems at 8 years

2c. Examiner-reported attention skills at 6 years

2d. Examiner-reported attention skills at 8 years

Legend:

 Regression line

 95% confidence intervals




Figure 1.




Figure 2.

Table 1. Description of the total sample and for appropriate for gestational age (AGA) and small for gestational age (SGA) children separately.

<b>Variables</b>		<b>Total (N=1435)</b>	<b>AGA (N=1059)</b>	<b>SGA (N=376)</b>	<b>p-values<sup>a</sup></b>
Gestational age in weeks	<i>Mean (SD)</i>	36.27 (4.11)	36.66 (4.15)	35.19 (3.78)	<0.001
Birth weight in grams	<i>Mean (SD)</i>	2582.14 (970.42)	2853.58 (914.68)	1817.64 (668.32)	<0.001
Prenatal complications	<i>Mean (SD)</i>	1.26 (1.21)	1.18 (1.17)	1.48 (1.29)	<0.001
Preeclampsia	<i>N (%)</i>	38 (2.7)	10 (0.9)	28 (7.5)	<0.001
Anemia	<i>N (%)</i>	9 (0.6)	3 (0.3)	6 (1.6)	<0.01
Urinary tract infection	<i>N (%)</i>	27 (1.9)	21 (2.0)	6 (1.6)	<i>Ns</i>
Bleeding before 28 weeks	<i>N (%)</i>	106 (7.4)	75 (7.1)	31 (8.2)	<i>Ns</i>
Bleeding after 28 weeks	<i>N (%)</i>	57 (4.0)	40 (3.8)	17 (4.5)	<i>Ns</i>
Pathologic CTG	<i>N (%)</i>	63 (4.4)	31 (2.9)	32 (8.5)	<0.001
Preterm labour	<i>N (%)</i>	401 (27.9)	24 (2.3)	127 (33.8)	<0.001
Multiples	<i>N (%)</i>	132 (9.2)	95 (9.0)	37 (9.9)	<i>Ns</i>
Nicotine addiction	<i>N (%)</i>	194 (13.5)	136 (12.8)	58 (15.4)	<i>Ns</i>
First booking after 12 weeks	<i>N (%)</i>	224 (15.6)	173 (16.3)	51 (13.6)	<i>Ns</i>
No regular check-ups	<i>N (%)</i>	25 (1.7)	16 (1.5)	9 (2.4)	<i>Ns</i>
Infectious disease	<i>N (%)</i>	197 (13.7)	142 (13.4)	55 (14.6)	<i>Ns</i>
Severe illness or accident	<i>N (%)</i>	263 (18.3)	188 (17.8)	75 (19.9)	<i>Ns</i>
(Oligo)-hydramnion	<i>N (%)</i>	74 (5.2)	46 (4.3)	26 (6.9)	<i>Ns</i>
Mode of delivery <sup>b</sup>					<0.001
Vaginal birth	<i>N (%)</i>	732 (51.0)	596 (56.3)	136 (36.2)	
Emergency caesarean birth	<i>N (%)</i>	124 (8.6)	88 (8.3)	36 (9.6)	
Elective caesarean birth	<i>N (%)</i>	313 (21.8)	178 (16.8)	135 (35.9)	
Head circumference <sup>c</sup> in cm	<i>Mean (SD)</i>	42.55 (1.45)	42.78 (1.36)	41.88 (1.52)	<0.001
Female	<i>N (%)</i>	699 (48.7)	515 (48.6)	184 (48.9)	<i>Ns</i>
Family SES					<0.05
SES – high	<i>N (%)</i>	423 (30.1)	331 (31.3)	92 (24.5)	
SES – middle	<i>N (%)</i>	543 (37.8)	397 (37.5)	146 (38.8)	
SES – low	<i>N (%)</i>	467 (32.5)	330 (31.2)	137 (36.4)	
Outcome variables					

Parent-reported attention problems at 6 years	<i>Mean (SD)</i>	3.49 (2.79)	3.30 (2.75)	4.02 (2.84)	<0.001
Parent-reported attention problems at 8 years	<i>Mean (SD)</i>	2.34 (2.41)	2.21 (2.35)	2.70 (2.56)	<0.001
Examiner-reported attention skills at 6 years	<i>Mean (SD)</i>	5.90 (1.58)	6.03 (1.56)	5.54 (1.57)	<0.001
Examiner-reported attention skills at 8 years	<i>Mean (SD)</i>	6.65 (1.28)	6.75 (1.26)	6.37 (1.31)	<0.001

<sup>a</sup> Variables were compared between AGA and SGA groups using One-Way ANOVA for continuous variables and  $\chi^2$  test for categorical variables. <sup>b</sup> Missing data: n=142 (9.9%) did not have elective caesarean birth but not clear whether delivery was via vaginal birth or emergency caesarean, n=124 (8.6%) did not have mode of delivery information. <sup>c</sup> Head circumference was measured at 5 months (corrected for prematurity).

Table 2. Standardized regression coefficients (95% confidence intervals) for parent-reported attention problems at 6 and 8 years.

Variables	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6
<b>Parent-reported attention problems at 6 years</b>						
Gestational age	-0.191*** (-0.241;-0.140)	-0.071 (-0.147;0.005)	-0.051 (-0.136;0.034)	0-.026 (-0.112;0.059)	-0.032 (-0.117;0.053)	-0.022 (-0.106;0.056)
Gestational age – quadratic term		.161*** (0.085;0.236)	0.165*** (0.088;0.241)	0.188*** (0.111;0.265)	0.167*** (0.089;0.244)	0.149** (0.072;0.225)
Prenatal complications <sup>a</sup>			0.042 (-0.015;0.098)	0.039 (-0.017;0.095)	0.030 (-0.026;0.087)	0.030 (-0.025;0.086)
Elective caesarean birth <sup>b</sup>			-0.003 (-0.058;0.051)	-0.022 (-0.077;0.033)	-0.013 (-0.068;0.041)	-0.007 (-0.061;0.047)
SGA <sup>c</sup>				0.108*** (0.055;0.161)	0.078** (0.024;0.133)	0.061** (0.007;0.115)
SGA*Gestational age				-0.026 (-0.082; 0.026)	-0.021 (-0.075; 0.032)	-0.009 (-0.063; 0.043)
Head Circumference <sup>d</sup>					-0.110*** (-0.163;-0.057)	-0.173*** (-0.230;-0.116)
Sex – FEMALE <sup>e</sup>						-0.169*** (-0.222; -0.115)
SES high <sup>f</sup>						-0.010 (-0.066;0.045)
SES low <sup>f</sup>						0.042 (-0.013;0.098)
R square <sup>g</sup>	0.04	0.05	0.05	0.06	0.07	0.10
<b>Parent-reported attention problems at 8 years</b>						
Gestational age	-0.154*** (-0.205;-0.103)	0.003 (-0.073;0.079)	0.009 (-0.076;0.094)	0.030 (-0.056;0.115)	0.024 (-0.061;0.110)	0.034 (-0.050;0.117)
Gestational age – quadratic term		0.211*** (0.135;0.287)	0.212*** (0.135;0.289)	0.232*** (0.155;0.310)	0.210** (0.133;0.288)	0.187*** (0.110;0.283)
Prenatal complications <sup>a</sup>			0.012 (-0.044;0.069)	0.010 (-0.046;0.067)	0.001 (-0.055;0.057)	0.006 (-0.049;0.062)
Elective			-0.001	-0.018	-0.009	-0.003

caesarean birth <sup>b</sup>			(-0.054;0.054)	(-0.073;0.037)	(-0.064;0.046)	(-0.057;0.052)
SGA <sup>c</sup>				0.093***	0.063*	0.042
				(0.039;0.146)	(0.008;0.118)	(-0.012;0.097)
SGA*Gestational				-0.024	-0.019	-0.008
age				(-0.080; 0.029)	(-0.74; 0.034)	(-0.061; 0.045)
Head					-0.110***	-0.184***
Circumference <sup>d</sup>					(-0.164;-0.057)	(-0.241;-0.127)
Sex- FEMALE <sup>e</sup>						-0.196***
						(-0.250; -0.143)
SES high <sup>f</sup>						-0.051
						(-0.107;0.004)
SES low <sup>f</sup>						-0.015
						(-0.070;0.041)
R square <sup>g</sup>	0.02	0.04	0.04	0.05	0.06	0.10

\* $p < 0.05$  \*\* $p < 0.01$  \*\*\* $p < 0.001$  <sup>a</sup> Higher scores on prenatal complications indicate more complications during pregnancy; <sup>b</sup> elective caesarean birth is a binary variable indicating elective caesarean birth versus delivery following labour; <sup>c</sup> SGA (small for gestational age) is a binary variable <sup>d</sup> Head circumference was measured at 5 months (corrected for prematurity) and higher scores in head circumference indicate bigger head circumference; <sup>e</sup> sex is a categorical variable (male = 0, female = 1); <sup>f</sup> family SES was included as two binary variables, SES-high and SES – low; <sup>g</sup> R square changes were significant ( $p < .001$ ) except for Model 3 (not significantly different from Model 2) and Model 6 (not significantly different from Model 5).

Table 3. Standardized regression coefficients (95% confidence intervals) for examiner-reported attention skills at 6 and 8 years.

Variables	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6
<b>Examiner-reported attention skills at 6 years</b>						
Gestational age	0.276*** (0.226;0.326)	0.143** (0.069;0.217)	0.120** (0.037;0.202)	0.093* (0.010;0.176)	0.098** (0.015;0.181)	0.091** (0.011;0.172)
Gestational age- quadratic term		-0.178*** (-0.252;-0.104)	-0.194** (-0.259;-0.109)	-0.210*** (-0.285;-0.134)	-0.191*** (-0.266;-0.115)	-0.168*** (-0.241;-0.094)
Prenatal complications <sup>a</sup>			-0.043 (-0.098;0.012)	-0.040 (-0.095;0.014)	-0.033 (-0.088;0.022)	-0.029 (-0.083;0.024)
Elective caesarean birth <sup>b</sup>			-0.002 (-0.055;0.051)	0.019 (-0.034;0.073)	0.011 (-0.042;0.065)	0.012 (-0.040;0.064)
SGA <sup>c</sup>				-0.118*** (-0.170;-0.066)	-0.092*** (-0.146;-0.039)	-0.071*** (-0.123;-0.019)
SGA*Gestational age				0.033 (-0.018; 0.087)	0.028 (-0.023; 0.082)	0.017 (-0.033; 0.068)
Head Circumference <sup>d</sup>					0.094*** (0.042;0.146)	0.139** (0.084;0.193)
Sex – FEMALE <sup>e</sup>						0.161*** (0.116; 0.222)
SES high <sup>f</sup>						0.169*** (0.116;0.222)
SES low <sup>f</sup>						-0.041 (-0.095;0.012)
R square <sup>g</sup>	0.08	0.09	0.09	0.11	0.12	0.18
<b>Examiner-reported attention skills at 8 years</b>						
Gestational age	0.255*** (0.205;0.305)	0.129** (0.055;0.204)	0.101* (0.018;0.184)	0.080 (-0.004;0.164)	0.088* (0.005;0.171)	0.079* (0.001;0.159)
Gestational age- quadratic term		-0.169** (-0.243;-0.094)	-0.174*** (-0.249;-0.098)	-0.197*** (-0.273;-0.121)	-0.167*** (-0.243;-0.091)	-0.142*** (-0.215;-0.059)
Prenatal complications <sup>a</sup>			-0.063* (-0.119;-0.008)	-0.061* (-0.116;-0.006)	-0.049 (-0.104;0.006)	-0.045(- 0.098;0.008)
Elective caesarean birth <sup>b</sup>			0.009 (-0.044;0.063)	0.029 (-0.025;0.083)	0.017 (-0.036;0.071)	0.016 (-0.036;0.067)

SGA <sup>c</sup>				-0.100***	-0.060*	-0.037*
				(-0.152;-0.048)	(-0.113;-0.007)	(-0.088;-0.015)
SGA*Gestational age				0.050	0.043	0.031
				(-0.001; 0.106)	(-0.008; 0.097)	(-0.020; 0.081)
Head Circumference <sup>d</sup>					0.149***	0.207***
					(0.097;0.200)	(0.153;0.261)
Sex- FEMALE <sup>e</sup>						0.192***
						(0.141; 0.243)
SES high <sup>f</sup>						0.141***
						(0.088;0.194)
SES low <sup>f</sup>						-0.061*
						(-0.114;-0.008)
R square <sup>g</sup>	0.07	0.08	0.08	0.09	0.11	0.18

\* $p < 0.05$  \*\* $p < 0.01$  \*\*\* $p < 0.001$  <sup>a</sup> Higher scores on prenatal complications indicate more complications during pregnancy; <sup>b</sup> elective caesarean birth is a binary variable indicating elective caesarean birth versus delivery following labour; <sup>c</sup> SGA (small for gestational age) is a binary variable <sup>d</sup> Head circumference was measured at 5 months (corrected for prematurity) and higher scores in head circumference indicate bigger head circumference; <sup>e</sup> sex is a categorical variable (male = 0, female = 1); <sup>f</sup> family SES was included as two binary variables, SES-high and SES – low; <sup>g</sup> R square changes were significant ( $p < .001$ ) except for Model 3 (not significantly different from Model 2) and Model 6 (not significantly different from Model 5).