



Identification of Aminoglycoside and β -Lactam Resistance Genes from within an Infant Gut Functional Metagenomic Library

Fiona Fouhy^{1,2}, Lesley A. Ogilvie^{3*}, Brian V. Jones^{3,4}, R. Paul Ross^{1,5}, Anthony C. Ryan⁶, Eugene M. Dempsey^{6,7}, Gerald F. Fitzgerald^{2,5}, Catherine Stanton^{1,5}, Paul D. Cotter^{1,5*}

1 Teagasc Food Research Centre, Moorepark, Fermoy, Cork, Ireland, **2** School of Microbiology, University College Cork, Cork, Ireland, **3** School of Pharmacy and Biomolecular Sciences, University of Brighton, Brighton, East Sussex, United Kingdom, **4** Queen Victoria Hospital NHS Foundation Trust, East Grinstead, West Sussex, United Kingdom, **5** Alimentary Pharmabiotic Centre, Cork, Ireland, **6** Department of Paediatrics and Child Health, University College Cork, Cork, Ireland, **7** Department of Neonatology, Cork University Maternity Hospital, Cork, Ireland

Abstract

The infant gut microbiota develops rapidly during the first 2 years of life, acquiring microorganisms from diverse sources. During this time, significant opportunities exist for the infant to acquire antibiotic resistant bacteria, which can become established and constitute the infant gut resistome. With increased antibiotic resistance limiting our ability to treat bacterial infections, investigations into resistance reservoirs are highly pertinent. This study aimed to explore the nascent resistome in antibiotic-naïve infant gut microbiomes, using a combination of metagenomic approaches. Faecal samples from 22 six-month-old infants without previous antibiotic exposure were used to construct a pooled metagenomic library, which was functionally screened for ampicillin and gentamicin resistance. Our library of ~220Mb contained 0.45 ampicillin resistant hits/Mb and 0.059 gentamicin resistant hits/Mb. PCR-based analysis of fosmid clones and uncloned metagenomic DNA, revealed a diverse and abundant aminoglycoside and β -lactam resistance reservoir within the infant gut, with resistance determinants exhibiting homology to those found in common gut inhabitants, including *Escherichia coli*, *Enterococcus* sp., and *Clostridium difficile*, as well as to genes from cryptic environmental bacteria. Notably, the genes identified differed from those revealed when a sequence-driven PCR-based screen of metagenomic DNA was employed. Carriage of these antibiotic resistance determinants conferred substantial, but varied (2–512x), increases in antibiotic resistance to their bacterial host. These data provide insights into the infant gut resistome, revealing the presence of a varied aminoglycoside and β -lactam resistance reservoir even in the absence of selective pressure, confirming the infant resistome establishes early in life, perhaps even at birth.

Citation: Fouhy F, Ogilvie LA, Jones BV, Ross RP, Ryan AC, et al. (2014) Identification of Aminoglycoside and β -Lactam Resistance Genes from within an Infant Gut Functional Metagenomic Library. PLoS ONE 9(9): e108016. doi:10.1371/journal.pone.0108016

Editor: Zaid Abdo, Agricultural Research Service, United States of America

Received: April 11, 2014; **Accepted:** August 21, 2014; **Published:** September 23, 2014

Copyright: © 2014 Fouhy et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. Full sequencing data sets available at <http://dx.doi.org/10.6070/H46971H6>.

Funding: FF is in receipt of an Irish Research Council EMBARK scholarship and is a Teagasc Walsh fellow. Research in the PDC laboratory is supported by the Irish Government under the National Development Plan through the Science Foundation Ireland Investigator award 11/PI/1137. This research was supported by the Science Foundation of Ireland-funded centre for Science Engineering and Technology, the Alimentary Pharmabiotic Centre, and the Irish Government under the National Development plan (INFANTMET: <http://eldermet.ucc.ie>). LAO and BVJ are funded by the UK Medical Research Council (grant number G0901553 awarded to BVJ), with work in the laboratory of BVJ also supported by the Hospital Infection Society, the Society for Applied Microbiology and the University of Brighton. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: PDC is a PLOS ONE editorial board member but this has not altered the authors' adherence to PLOS ONE editorial policies and criteria.

* Email: paul.cotter@teagasc.ie

‡ Current address: Alacris Theranostics GmbH, Berlin, Germany

Introduction

There is growing concern that we are rapidly approaching a post-antibiotic era. As a result every effort is being made to discover and investigate antibiotic resistance reservoirs with the aim of limiting the selection for, or dissemination of, antibiotic resistance genes. One such reservoir is the human gut microbiota. Colonized by trillions of bacteria representing hundreds of different species, this ecosystem has been identified as a source of antibiotic resistant bacteria [1–3]. Though vastly less complex and considerably less stable than the adult gut microbiota, the gut microbiota of infants also has the potential to acquire and

disseminate antibiotic resistant genes [4]. From the commencement of labour, the sterile infant gut rapidly becomes colonized [5] and from birth through to 2 years of age the infant gut microbiota is dynamic, unstable and becomes increasingly complex, until it resembles that of an adult. The identity of the first colonizers of the infant gut depends on numerous factors [6,7] including mode of delivery [8], and pre-term versus full-term gestation [9] and, because of the aforementioned instability of this microbial population, many factors including feeding choice [10], probiotic or prebiotic supplementation [11] and antibiotic exposure [12] can significantly impact on its subsequent development. Indeed, in a previous study, we have demonstrated that the infant microbiota

becomes dominated by *Proteobacteria* following ampicillin and gentamicin administration in the first 48 hours of life [13], which may be due to the known high prevalence of antibiotic resistant species within this phylum [14]. Such findings suggest that the infant gut microbiota, though immature and in constant flux, can be a source of resistant bacteria which can become dominant following antibiotic exposure. Given the instability of the infant microbiota in early life, there is considerable opportunity for the infant gut to acquire resistant populations which, if they become established, could have significant effects on shaping the composition of the microbiota later in life [15].

While it is known that antibiotic resistance genes are present within the gut microbiota from early life [4,16], data with respect to the presence of antibiotic resistance genes in healthy infants with no history of antibiotic treatment is still limited [4]. Furthermore, much of the existing data comes from studies investigating resistance genes in specific commensals, such as strains of *Escherichia coli* or *Lactobacillus* [17,18], rather than the entire microbiota. Challenges in studying gut microbiota are well documented [6] and include an inability to culture the majority of gut microbes in a laboratory environment [19]. However, metagenomic libraries provide the opportunity to capture metagenomic DNA from complex environments and to functionally screen for phenotypes of interest. The limited existing data available from adult [1,20] and infant [21,22] metagenomic banks indicate that human gut microbiota is a source of diverse antibiotic resistance genes. To date, however, there is a significant paucity in infant functional metagenomic studies, particularly relating to infants free from antibiotic treatment. Thus, the aim of this study was to construct a fosmid bank using metagenomic DNA from 6-month-old infants who had never received antibiotic treatment and to screen for the presence of antibiotic resistance genes. By 6 months of age, the infants will have acquired an increasingly complex microbiota and will have had the opportunity to acquire antibiotic resistant strains from numerous sources [5,23]. Our goal was to provide insight into the infant gut microbiota as a reservoir for resistance genes, when no antibiotic selective pressure exists, using a combination of functional metagenomics and PCR-based approaches. Due to the on-going focus of our research [13], we specifically investigated resistance to aminoglycosides and β -lactam antibiotics. Using both a function-based and function-independent approach, we successfully identified a variety of aminoglycoside and β -lactam resistance genes in the gut microbiota of infants, providing insights into the infant gut resistome.

Materials and Methods

Recruitment of volunteers

Infants were recruited as part of the INFANTMET study (<http://eldermet.ucc.ie>). Parents of infant participants provided written informed consent. Approval for the INFANTMET study was received from the Clinical Research Ethics Committee of the Cork Teaching Hospitals, Cork, Ireland. Infants participating in the study had no antibiotics administered for the first 6 months of life. Fresh faecal samples were collected from 22 6-month-old infants and immediately stored at -80°C until processed.

DNA extraction

Due to the small volume of each individual sample, it was necessary to pool faecal samples prior to DNA extraction. Faecal samples were homogenized and a 500 mg aliquot from each were pooled to form one sample from which high molecular weight

metagenomic DNA was extracted, using a previously described method [24–26]. Briefly, faecal samples were homogenized in PBS (Sigma Aldrich, Dublin, Ireland), centrifuged at $1000\text{g}\times 5\text{mins}$ and the supernatant retained. Nycodenz (Axis Shield, UK) density gradient separation was performed to separate out the bacterial cells. Enzymatic digestion of the cells using lysozyme and mutanolysin (Sigma Aldrich) was performed, and subsequently the protein was removed using Proteinase K and ammonium acetate treatment (Sigma Aldrich). DNA was then purified and precipitated using standard chloroform and ethanol precipitation procedures and was eluted in TE buffer.

Metagenomic bank creation

To allow functional screening of the metagenomic DNA from the infant gut, a fosmid metagenomic bank was created using the EpiCentre CopyControl Fosmid Library Production kit with the pCC1FOS vector, according to the manufacturer's instructions (Cambio, Cambridge, England). Briefly, size selection was performed on the metagenomic DNA using pulse field gel electrophoresis (PFGE) ($0.5\times \text{TAE}$; 0.1 initial/10 final switch times; 4V; 17 hours; 14°C) and fragments of $\sim 40\text{ Kb}$ were gel extracted from the low melting point agarose (Promega, Medical Supply Company, Dublin). Fragments were then ligated with the pCC1FOS vector according to the manufacturer's instructions and subsequently packaged into EPI300-T1^R *E. coli* plating strain cells. These cells were then plated onto LB agar plates (Difco, Becton, Dickinson & Co, Oxford, England) containing $12.5\ \mu\text{g/ml}$ chloramphenicol, IPTG and Xgal (Sigma Aldrich) and grown overnight aerobically at 37°C . To verify the diversity of the library, random white clones were selected and were digested with *Pst*I and *Nde*I restriction enzymes (New England Biolabs, UK), to determine if different DNA insert sequences were present in our metagenomic bank. The entire library was plated and then picked and stocked in 384-well-format using the QPix2-XT robotic system (Molecular Devices, Berkshire, UK) and was stored at -80°C until screening. A library of $\sim 220\text{Mb}$ of DNA was created.

Screening for antibiotic resistant clones

Due to the ongoing focus of our research [13,27], we chose to concentrate on the resistance of infant gut microbiota to 2 groups of antibiotics, namely the aminoglycosides and β -lactams. For screening of the metagenomic bank, the library was plated onto LB agar with $12.5\ \mu\text{g/ml}$ chloramphenicol and inhibitory concentrations of ampicillin ($50\ \mu\text{g/ml}$) or gentamicin ($10\ \mu\text{g/ml}$) (Sigma Aldrich). These levels of antibiotics were chosen due to their inhibition of host EPI 300-T1^R *E. coli* cells. The library was plated in triplicate and plates were incubated aerobically at 37°C for 24–36 hours. Clones found to be resistant over three replicate plates were selected, their resistance phenotype verified by re-streaking onto agar containing the relevant antibiotic and were stocked for further analysis.

PCR analysis

To investigate which resistance genes were present in the insert DNA that conferred resistance, PCRs were carried out on each of these clones using primers for aminoglycoside and β -lactam resistance genes (Table 1). For the aminoglycosides, we used degenerate primers designed to amplify the acetylation (AAC; *aac* (3)-I, *aac* (3)-II, *aac* (3)-III, *aac* (3)-VI and *aac* (6)), adenylation (ANT; *ant* (2'')-Ia) and phosphorylation (APH; *aph* (2'')-Ic and *aph* (2'')-Id) genes [28], as well as the bifunctional gene *aac* (6')-Ie-*aph*(2'')-Ia [29,30]. For β -lactam resistance the primer sets for the

following genes were used *bla*_{TEM} [31,32], *bla*_{OXA} [33], *bla*_{SHV} [33], *bla*_{ROB} [32] and *bla*_{CTX-M} [34] (Table 1). Resistant clones were grown overnight in LB broth supplemented with chloramphenicol (12.5 µg/ml) and either ampicillin (50 µg/ml) or gentamicin (10 µg/ml). Fosmids were extracted from the clones using the QIAprep Spin Mini Prep kit (Qiagen, Sussex, UK) and subsequently used as template DNA for PCR analysis. PCRs were performed using previously outlined protocols [28–34]. Each reaction contained 25 µl of Biomix Red (MyBio, UK), 1 µl forward primer (10pmol), 1 µl reverse primer (10pmol), fosmid DNA (at the volume required to correspond to 64 ng) from the resistant clone, and PCR grade water (Bioline, Medical Supply Company, Dublin, Ireland) to a final reaction volume of 50 µl. All reactions were performed in duplicate. PCR products were visualized using gel electrophoresis (1.5% agarose, 1× TAE, 100V). Negative controls, in which the DNA template was substituted with PCR-grade water, failed to generate an amplicon, ruling out the possibility of contamination. Successful duplicate PCRs were pooled and cleaned using AMPure (Beckman Coulter UK) magnetic bead-based purification procedures.

To determine if additional resistance genes were present in the infant gut microbiome that had not been captured in the metagenomic bank, additional PCRs for detecting resistance genes were performed on the metagenomic DNA which was used for construction of the fosmid bank. PCR analysis was performed using the primers and protocols outlined above. Following PCR amplification, amplicons were cloned using the TOPO TA cloning kit (Invitrogen, Dublin, Ireland) according to the manufacturer's instructions. TOPO cloning reactions were then transformed into *E. coli* TOP 10 cells and plated on LB agar containing antibiotics for the selection of the cloning vector (either kanamycin 50 µg/ml or ampicillin 50 µg/ml). Plasmids were then extracted from overnight cultures of the TOPO sub-clones using the QIAprep Spin Mini Prep kit (Qiagen, Sussex, UK) to facilitate subsequent DNA sequencing.

Sequencing and analysis

A subset of PCR products from the resistant fosmid clones were sent for Sanger sequencing to determine their closest homologue (Source Biosciences, Dublin, Ireland). The subset included representatives from each of the genes investigated and from clones containing multiple resistance genes. Plasmid DNA from a subset of the TOPO sub-clones of the metagenomic DNA PCR products were also sent for Sanger sequencing. Sequencing reads were BLASTed against the NCBI non-redundant database using BLASTx (<http://blast.ncbi.nlm.nih.gov/>). In the event where multiple hits occurred, the BLAST hit which displayed greatest homology (based on E value) is reported.

Minimum inhibitory concentration assays

Minimum inhibitory concentration (MIC) tests were conducted on gentamicin (Gent^R) and ampicillin resistant (Amp^R) fosmid clones to determine the level of resistance conferred by the insert DNA. In order to determine the relative increase in resistance, MICs were compared to the control strain (EPI300-T1^R *E. coli* + empty pCC1FOS fosmid). MICs were performed according to the British Society for Antimicrobial Chemotherapy guidelines [35]. An antibiotic range of 256–0.25 mg/L was used. Positive (broth + inoculum only) and negative controls (broth + antibiotic only) were included in each assay. Plates were incubated aerobically at 37°C for 24 hours. The MIC was determined as the lowest concentration of antibiotic at which no visible growth occurred.

Results

Identification of gentamicin and ampicillin resistant clones within the infant gut microbiome using a functional metagenomic bank

Ten white clones were picked at random from LB agar plates containing 12.5 µg/ml chloramphenicol, IPTG and X-gal and underwent restriction digestion using *Pst*I and *Nde*I restriction enzymes. These randomly selected clones had unique restriction profiles (data not shown), establishing that the metagenomic library contained different insert DNA, demonstrating a diverse metagenomic fosmid bank had been successfully created. Our library contained ~220Mb of metagenomic DNA, which contained 0.45 ampicillin resistant hits/Mb and 0.059 gentamicin resistant hits/Mb. To determine which genes were responsible for the resistance in the isolated clones, PCR analysis was conducted on each of the resistant clones using primers for β-lactam and aminoglycoside resistance genes (sequencing files available at <http://dx.doi.org/10.6070/H46971H6>). Furthermore, to determine if additional genes could be detected in the uncloned metagenomic DNA, PCR analysis was completed on this DNA with the same degenerate primers as used on the resistant clones (Figure 1).

PCR and Sanger sequencing of β-lactam resistant clones

Analysis of the one hundred Amp^R clones using *bla*_{TEM} primers, revealed that they all contained *bla*_{TEM} genes (Table 2). Based on sequencing analysis, these genes shared closest homology with *bla*_{TEM} genes from uncultured soil bacteria (clones 2, 3, 4, 5, 6, 10, 16, 17, 123, 128), *Shigella* sp. (clone 4), *Serratia marcescens* (clone 126) and *Citrobacter freundii* (clones 17 and 18) (Table 2). Sequencing of amplicons generated using the same *bla*_{TEM} primers and uncloned metagenomic DNA, revealed a number of different *bla*_{TEM} genes that shared closest homology with genes from *Klebsiella pneumoniae*, *E. coli*, *Salmonella enterica* and *Staphylococcus aureus* (Table 3).

PCRs revealed that half of the Amp^R clones contained *bla*_{CTX-M} genes (including CTX-M-1 and CTX-M-36 gene homologues), which shared homology with 1 of 3 different *bla*_{CTX-M} genes present in *E. coli*. Sequencing of *bla*_{CTX-M} PCR products generated directly from metagenomic DNA identified additional sources of these genes, detecting genes that shared homology with *bla*_{CTX-M} genes present in *E. coli* (including one source that was not detected in the analysis of the Amp^R clones) and *K. pneumoniae* (Table 3).

Just 2 of the 100 Amp^R clones gave PCR products using the *bla*_{OXA} primers, which surprisingly resembled 2 different *bla*_{TEM} genes, rather than *bla*_{OXA} genes, from a variety of *E. coli*. No *bla*_{OXA} genes were amplified when PCR was directly applied to the metagenomic DNA from the infant faecal samples. Six of the 100 Amp^R clones (clones 15, 50, 51, 61, 62 and 63) contained *bla*_{ROB} genes and all shared closest homology (53% identity, 40% query cover) with a hypothetical protein from *Staphylococcus haemolyticus*. Direct PCR amplification using template metagenomic DNA failed to reveal any *bla*_{ROB} genes. Finally, *bla*_{SHV} primers did not generate amplicons from either the Amp^R clones or uncloned metagenomic DNA.

PCR and Sanger sequencing of aminoglycoside resistant clones

Degenerate PCR primers were employed to determine whether the fosmids conferring gentamicin resistance contained homologues of known aminoglycoside resistance genes. Acetylation, adenylation and phosphorylation genes were detected in the

Table 1. Primers used in this study.

Target gene family	Primer name	Sequence 5'-3'	Annealing temperature °C	Ref.
β-lactamase genes				
<i>bla</i> _{TEM}	RH605	TTTCGTGTCGCCCTTATTCC	60	Bailey <i>et al.</i> (2011)
	RH606	CCGGCTCCAGATTATCAGC		
	Bla_TEMF	TGGGTGCACGAGTGGTTAC	57	Tenover <i>et al.</i> (1994)
	Bla_TEMR	TTATCCGCCTCCATCCAGTC		
<i>bla</i> _{ROB}	Bla_ROBF	ATCAGCCACACAAGCCACCT	62	Tenover <i>et al.</i> (1994)
	Bla_ROBR	GTTTGCGATTGGTATGCGA		
<i>bla</i> _{SHV}	Bla_SHVF	CACTCAAGGATGTATTGTG	58	Briñas <i>et al.</i> (2002)
	Bla_SHVR	TTAGCGTTGCCAGTGCTCG		
<i>bla</i> _{OXA}	Bla_OXAF	TTCAAGCAAAGGCACGATAG	64	Briñas <i>et al.</i> (2002)
	Bla_OXAR	TCCGAGTTGACTGCCGGTTG		
<i>bla</i> _{CTX-M}	Bla_CTX-MF	CGTTGTAAAACGACGGCCAGTGAATGT-GCAGYACCAGTAARGTKATGGC	55	Monstein <i>et al.</i> (2009)
	Bla_CTX-MR	TGGGTRAARTARGTSACCAGAAYCAGCGG	60	
AG resistance genes				
<i>aac</i> (3)-I	Faac3-1	TTCATCGCGCTTGCTGCTTYTYGA	58	Heuer <i>et al.</i> (2002)
	Raac3-1	GCCACTGCGGATCGTRCCRTA		
<i>aac</i> (3)-II/VI	Faac3-2	GCGCACCCGATGCMTCATGG	58	
	Raac3-2	GGCAACGGCCTCGGCTARTGSA		
	Faac3-6	GCCCATCCGACGCATCSATGG		
	Raac3-6	CGCCACCGCTTCGGCATARTGSA		
<i>aac</i> (6')-II/Ib	Faac6	CACAGTCGTACGTTGCKTBGG	58	
	Raac6	CCTGCCTTCTCGTAGCAKCGDAT		
<i>ant</i> (2'')-Ia	Fant	TGGGCGATCGATGCACGGCTRG	58	
	Rant	AAAGCGGCACGCAAGACCTCMAC		
<i>aph</i> (2'')-I	Faphc	CCCAAGAGTCAACAAGGTGCAGA	55	
	Faphd	GGCAATGACTGTATTGCATATGA	55	
	Raph	GAATCTCCAAAATCRATWATKCC		
<i>aac</i> (6')-Ie- <i>aph</i> (2'')-Ia	aac-aphF	GAGCAATAAGGGCATAACAAAATC	47	De Fatima Silva Lopes <i>et al.</i> (2003)
	aac-aphR	CCGTGCATTTGTCTTAAAAACTGG		
	aac6-aph2F	CCAAGAGCAATAAGGGCATAACC	55	Schmitz <i>et al.</i> (1999)
	aac6-aph2R	CACACTATCATAACCATCACCG		

AG: aminoglycoside.

doi:10.1371/journal.pone.0108016.t001

Gent^R clones (Table 2). With respect to aminoglycoside acetylation-associated genes, *aac* (3)-I, *aac* (3)-II, *aac* (3)-III and *aac* (3)-VI were not detected in either the 13 Gent^R clones or in the uncloned metagenomic DNA. However, all 13 Gent^R clones contained an *aac* (6) gene that most closely resembled a *Pseudomonas fluorescens* gene. Interestingly, the *aac* (6) primers did not generate amplicons when used directly with metagenomic DNA (Table 3). Aminoglycoside adenylation, *ant* (2'')-Ia, genes were detected in 9 of the 13 Gent^R clones (Table 2). Eight of these 9 Gent^R clones shared closest homology with aminoglycoside 2' O adenylation genes found in *Pasteurella multocida*, while that from the remaining clone resembled an *E. coli* associated gene. Similar sources of these genes were also detected when PCR was performed with the *ant* (2'')-Ia primers and uncloned metagenomic template DNA (Table 3). Following cloning and sequencing of these amplicons, it was apparent that 4 of the 5 clones contained *ant* (2'')-Ia genes sharing closest homology (99% identity, 97%

query cover) with genes present in *P. multocida* (identical to the source of these genes detected in the Gent^R clones). The fifth clone that was sequenced contained insert DNA sharing closest homology with a nucleotidyltransferase from *A. baumannii* (99% identity, 97% query cover).

When primers were applied to investigate the presence of aminoglycoside phosphorylation genes using *aph* (2'')-Id primers, 10 of the 13 Gent^R clones contained these genes (Table 2). A number of these Gent^R clones (5, 6, 8, 11, 12, 13 and 34) shared closest homology with phosphorylation genes from *Clostridium difficile* (100% identity and 94% query cover). When these *aph* (2'')-Id primers were used directly with metagenomic DNA, genes sharing homology with *C. difficile* (identical to those detected in the Gent^R clones), *Enterococcus faecium* and an unknown source of *aph* (2'')-Id genes were detected (Table 3).

Finally, using 2 different primer sets, it was found that all Gent^R clones contained the bifunctional gene *aac*(6')-Ie-*aph*(2'')-Ia. These




Figure 1. Schematic of the approach applied in this study.
doi:10.1371/journal.pone.0108016.g001

genes shared homology with bifunctional *aac(6')-Ie-aph(2'')-Ia* genes from *Enterococcus faecalis* (clones 3, 4, 5, 6 and 11), *Streptococcus epidermidis* (clones 4, 8 and 13), *E. faecium* (clone 10) and *Enterococcus* sp. (clone 5) (Table 2). Sequencing of the amplicons generated with uncloned metagenomic DNA, identified genes sharing closest homology with bifunctional genes from *E. faecalis* (which were distinct from those detected in the Gent^R clones) and from *S. aureus* (WP_001028140.1), a source which was not detected using PCR analysis on the Gent^R clones (Table 3).

MIC analysis of ampicillin and gentamicin resistant clones

To determine the level of resistance conferred by the fosmid-cloned metagenomic DNA to ampicillin or gentamicin, MIC assays were conducted. A representative subset of 22 of the 100 Amp^R clones (including those containing genes only detected in a limited number of clones e.g. *bla*_{OXA} and clones containing multiple β-lactam resistance genes) and all 13 of the Gent^R clones were studied. MICs were compared to the host EPI 300-T1^R *E. coli* cells containing the empty pCC1FOS fosmid.

The Amp^R clones exhibited a 2- (clone 6) to 512- (clone 2) fold increase in resistance to ampicillin compared to the empty fosmid control (Table 4). The large variation in MICs between the Amp^R clones highlighted the fact that different, but related, genes

conferred different levels of resistance to the host *E. coli* cells, most likely due to differences in expression/translation in the surrogate *E. coli* host. Clones containing multiple β-lactam resistance genes (clones 2, 3, 4, 62, 128, 136 and 140), including *bla*_{TEM}, *bla*_{CTX-M} and *bla*_{OXA} genes, were found to confer the highest levels of resistance to β-lactams. In contrast, when MICs were conducted on the 13 Gent^R clones, all clones showed similar levels of resistance to gentamicin with MICs of 4-8 fold higher than that of the control (Table 5).

Discussion

The infant gut microbiota is in constant flux during the first two years of life [23]. During this time there is considerable opportunity for the infant to acquire antibiotic resistant populations. Given the opportunity, resistant populations could become dominant in the gut following antibiotic therapy or may contribute to antibiotic resistance gene dissemination. In this study we set out to use a metagenomic fosmid bank to functionally screen the gut microbiota of 22 healthy infants who had not been treated with antibiotics, for antibiotic resistance genes. Additionally, we wanted to perform PCR analysis directly on uncloned metagenomic DNA to determine if additional genes would be detected, that had not been captured in the functional bank. Using such an approach, we

Table 2. Identity of ampicillin and gentamicin resistant genes amplified from the infant gut metagenomic bank.

Accession number	Primer name	Clones in which genes detected	Closest homologue*	E value	% ID
ROB					
	Bla_ROBF				
	Bla_ROBR				
WP_016930608.1		15,50,51,61,62,63	Hypothetical protein <i>S. haemolyticus</i>	1e ⁻²⁰	53
TEM					
	RH605				
	RH606				
	Bla_TEMF				
	Bla_TEMR				
AGH19654.1		2, 5,6	β-lactamase partial uncultured soil bacterium	1e ⁻¹³⁴	100
AGH19657.1		3,16,17	β-lactamase partial uncultured soil bacterium	4e ⁻¹³⁹	99
AAP93842.1		17,18	β-lactamase <i>C. freudii</i>	5e ⁻¹³⁸	99
AEN75339.1		4	β-lactamase TEM <i>Shigella</i> sp.	4e ⁻¹⁴¹	99
AGH19655.1		2,123	β-lactamase uncultured soil bacterium	2e ⁻⁹⁷	100
AGH19650.1		3,4,10,128	β-lactamase uncultured soil bacterium	7e ⁻⁹⁷	100
AAP93841.1		126	β-lactamase <i>S. marcescens</i>	2e ⁻⁹⁷	98
CTX-M					
	Bla_CTX-MF				
	Bla_CTX-MR				
AEZ49563.1		2,3	β-lactamase CTX-M-1 <i>E. coli</i>	2e ⁻¹¹⁶	99
BAD16611.1		4,136	β-lactamase CTX-M-36 <i>E. coli</i>	1.10e ⁻¹²⁷	99
AAB22638.1		140,145	β-lactamase penicillin amido β-lactamase hydrolase <i>E. coli</i>	2e ⁻¹²⁸	100
OXA					
	Bla_OXAF				
	Bla_OXAR				
AGN75112.1		140	TEM-190 β-lactamase <i>E. coli</i>	2e ⁻⁰⁴	38
ADZ11076.1		132	β-lactamase TEM <i>E. coli</i>	2e ⁻⁴⁰	71
ant (2'')-Ia					
	Fant				
	Rant				
YP_005176240.1		3,7,9,10,11,12,33,34,36	AG 2 O' adenylyltransferase <i>P. multocida</i>	1e ⁻⁷⁹	96
WP_000292466.1		8	AG adenylyltransferase <i>E. coli</i>	5e ⁻⁷⁵	96
aph (2'')-Id					
	Faphd				
	Raph				
WP_0214010241.1		5, 6, 8,11,12,13, 34,	aph <i>C. difficile</i>	4e ⁻⁹⁵	100
aac(6)					
	Faac6				
	Raac6				
AAA25680.1		34,36	AG 6' N acetyltransferase <i>P. fluorescens</i>	5e ⁻³³	97
aac (6')-Ie-aph (2'')-Ia					
	aac-aphF				
	aac-aphR				
	aac6-aph2F				
	aac6-aph2R				
WP_010714603.1		3,4,5,6,11	bifunctional aac (6')-Ie-aph (2'') <i>E. faecalis</i>	1e ⁻⁹⁶	100
AFR11868.1		4,8,13	bifunctional 6' AG N acetyltransferase/2'' AG phosphotransferase <i>S. epidermidis</i>	8e ⁻³²	98
AFM29914.1		5	Gentamicin resistance protein <i>Enterococcus</i> sp.	3e ⁻³⁰	96
WP_010782592.1		10	Bifunctional AAC/APH <i>E. faecium</i>	1e ⁻³¹	98

* Result presented is the closest homologue to our sequence (based on lowest E value).
doi:10.1371/journal.pone.0108016.t002

have demonstrated that the infant gut is a source of a variety of genes encoding resistance to aminoglycosides and β-lactams. One hundred Amp^R clones were detected in the gut microbiota of infants through screening of the metagenomic fosmid bank. Applying a PCR-based approach to determine the resistance genes

present in the resistant clones, we detected *bla*_{TEM}, *bla*_{OXA}, *bla*_{ROB} and *bla*_{CTX-M} genes. Interestingly, the *bla*_{TEM} genes detected exhibited greatest homology with genes from uncultured bacteria. Such results highlighted the ability of our strategy to reveal genes from more cryptic sources, including β-lactam

Table 3. Sequencing results from TOPO cloning of ampicillin and gentamicin resistant genes from metagenomic DNA.

Accession #	Closest homologue*	E value	% ID
TEM			
AAL03985.1	ESBL TEM-71 <i>K. pneumoniae</i>	7e ⁻¹⁵⁴	99
WP_004207849.1	β-lactamase TEM partial <i>K. pneumoniae</i>	3e ⁻¹⁵³	97
WP_017431996.1	β-lactamase partial <i>S. aureus</i>	1e ⁻¹¹²	84
WP_019405145.1	β-lactamase partial <i>K. pneumoniae</i>	6e ⁻¹⁵⁴	100
AEN02824.1	β-lactamase TEM-1 <i>K. pneumoniae</i>	4e ⁻¹¹¹	99
ADE18896.1	TEM-1 β-lactamase <i>S. enterica</i>	7e ⁻¹¹²	97
ABG46354.1	ESBL <i>E. coli</i>	1e ⁻¹³⁹	100
AEN02826.1	β-lactamase TEM-1 <i>K. pneumoniae</i>	1e ⁻¹⁰⁸	99
CTX-M			
AAB22638.1	β-lactamase penicillin amido β-lactam hydrolase <i>E. coli</i>	1e ⁻¹³⁹	100
AEZ49551.1	β-lactamase CTX-M-1 <i>K. pneumoniae</i>	5e ⁻¹²⁹	99
AEZ49563.1	β-lactamase CTX-M-1 <i>E. coli</i>	8e ⁻¹⁴⁰	97
ABG46356.1	ESBL <i>K. pneumoniae</i>	1e ⁻¹³⁸	99
ant (2'')-Ia			
YP_005176240.1	AG 2 O adenylyltransferase <i>Pasteurella multocida</i>	1e ⁻⁹⁵	99
WP_000946493.1	2 AG nucleotidyltransferase <i>A. baumannii</i>	3e ⁻⁹⁷	99
aph (2'')-Id			
AAW59417.1	<i>E. faecium</i> aph 2 Id	5e ⁻⁹⁸	90
3Sg8_9	Crystal structure of AG 2'' phosphotransferase Tobramycin resistance gene	5e ⁻¹⁰⁹	100
WP_021401024.1	aph (2'')-Id <i>C. difficile</i>	6e ⁻¹⁰⁸	98
aac (6')-Ie-aph (2'')-Ia			
WP_001028140.1	GNAT family acetyltransferase <i>S. aureus</i>	2e ⁻¹⁰⁹	100
AAX82584.1	Bifunctional AG modifying enzyme <i>E. faecalis</i>	2e ⁻¹⁰⁶	99
WP_002417297.1	Phosphotransferase enzyme family protein <i>E. faecalis</i>	4e ⁻¹¹³	99

* Result presented is the closest homologue to our sequence based on lowest E value.

ESBL: extended spectrum β-lactamase.

AG: aminoglycoside.

doi:10.1371/journal.pone.0108016.t003

resistance genes from sources not accessible through classical culture-based approaches. It was also notable that we detected such a prevalence of ampicillin resistance genes, and that they appear to have originated from a diverse range of species. Nonetheless, the majority of the β-lactam resistance genes identified shared homology with β-lactam resistance genes present in *E. coli*, *Shigella* and *Serratia*, which are common sources of such genes [36–38]. When PCRs were performed using *bla*_{OXA} primers, just 2 of the 100 Amp^R clones were found to contain such genes. However, sequencing of these amplicons revealed that they shared greatest homology with *bla*_{TEM} genes (Table 2). Therefore, it appears that in this instance, the infant gut microbiota did not contain *bla*_{OXA} genes and in their absence, the *bla*_{OXA} primers enabled the detection of *bla*_{TEM} homologues. This is not entirely unexpected, given previous observations of 3 sites of homology between the amino acid sequence of the OXA 2 gene and TEM β-lactamases [39].

This metagenomic study has successfully identified acetylation, adenylation and phosphorylation genes conferring aminoglycoside resistance in the infant gut microbiota. The fact that these genes resemble genes from bacteria such as *E. coli*, *E. faecalis*, *S. epidermidis* and *C. difficile* is not surprising given that a number of the observed resistance genes are resident in such species [30,40,41]. A recent study which employed a metagenomic

approach to investigate antibiotic resistance in infants, also detected a high prevalence of aminoglycoside resistance genes [22]. In that instance, they too identified aminoglycoside acetylation, adenylation and phosphorylation genes in the infant gut which shared homology with *C. difficile* and *E. faecium*. As these bacteria may be among the first colonizers of the infant gut, these findings support previous research which suggests antibiotic resistance and the antibiotic resistome is established in early life, perhaps even from birth, irrespective of antibiotic exposure, and is closely associated with microbes from the maternal and environmental sources it is exposed to during and immediately after birth [4]. This is consistent with the apparently ubiquitous nature of antibiotic resistance genes in the environment, with studies having demonstrated that antibiotic resistant isolates can be detected in individuals from remote areas of the world who lack, or who have minimal, antibiotic exposure [42–44].

In addition to screening the fosmid metagenomic bank, a PCR-based analysis of uncloned metagenomic DNA was also carried out. Previous studies demonstrated the success of such an approach with respect to detecting genes that were present at a low level and had not been captured in the metagenomic fosmid bank [21]. Additionally, this approach also addresses situations whereby certain genes may not be expressed or their products may not be functional in *E. coli*, e.g. as previously shown in the case of

Table 4. MICs of ampicillin resistant clones.

Amp ^R clone	MIC fold difference compared to the control
2	512
3	128
4	256
5	4
6	2
10	16
15	4
16	32
17	64
18	32
50	4
51	16
61	64
62	256
63	4
123	64
126	16
128	128
132	16
136	256
140	128
145	8

doi:10.1371/journal.pone.0108016.t004

strA and *strB* [45]. However, previous studies have demonstrated that genes from at least 4 major phyla from the gut microbiome can be functional in *E. coli* hosts [46]. It is also possible to create metagenomic banks using Gram-positive hosts such as *Streptomyces* to identify genes which may not be expressed in Gram-negative hosts [47]. In our study, as our library was only made in a Gram-negative host, we conducted PCR analysis directly on uncloned

Table 5. MICs of gentamicin resistant clones.

Gent ^R clones	MIC fold difference compared to the control
3	8
4	8
5	8
6	8
7	8
8	8
9	8
10	8
11	8
12	8
13	4
33	4
34	8

doi:10.1371/journal.pone.0108016.t005

metagenomic DNA, with the aim of identifying additional resistance genes to those captured in our functional screen. Thus, in this instance, we have demonstrated the usefulness of supplementing functional metagenomics with a PCR-based approach directly on metagenomic DNA to enable detection of antibiotic resistance genes apparently originating from different species than those that were identified during the course of the functional screen. In the majority of cases the putative sources of genes detected by PCR with metagenomic library isolates and with uncloned metagenomic template DNA were similar, though in some circumstances additional insights were provided using this combined approach. For example, some *bla*_{TEM} genes identified from the functional screen shared closest homology with genes from uncultured soil bacterium, whereas direct PCR-screening highlighted homologues of genes found in *K. pneumoniae*, *S. aureus*, *S. enterica* and *E. coli*. Furthermore, in certain instances resistance genes were captured by the metagenomic bank (*aac* 6) that were not detected when PCR analysis was completed directly on metagenomic DNA. This demonstrates that combining different experimental approaches allows for the most comprehensive insight into the gut microbiota as a source of antibiotic resistance genes. It should also be noted that, if screened for, we expect that our metagenomic library would also uncover resistance genes for other major families of antibiotics. Multi-drug resistant Gram-positive bacteria represent a serious threat to human health [48]. Several recent functional metagenomic studies screened the gut microbiota of children for resistance against antibiotics which specifically target Gram-positive bacteria (including fosfomycin and tigecycline) [20,22]. In the study of Moore *et al.*, clones resistant to antibiotics which target Gram-positive bacteria were detected [22], demonstrating that the gut microbiota of children and teenagers is a reservoir for resistant Gram-positive bacteria. Thus, while our focus was resistance to aminoglycoside and β -lactam antibiotics, future screening of this metagenomic bank for resistance to other antibiotics would likely yield positive clones.

This study is one of a limited number that has demonstrated the existence of antibiotic resistant genes in the healthy infant gut microbiota using a metagenomic library approach. Although used to investigate antibiotic resistance in environments as diverse as gull faeces [49] or soil from apple orchards [45], to date the use of metagenomic banks to screen the infant gut microbiota for resistance genes has been limited [21,22]. In particular, our study adds considerably to current data on the infant gut resistome by supporting and supplementing existing results [22], which have demonstrated that the infant gut is a reservoir for antibiotic resistance. By studying 6-month-old infants with no previous antibiotic exposure, we were able to establish a comprehensive insight into the resistome of infants devoid of antibiotic selective pressure. Our results supplement those of Moore *et al.* who investigated antibiotic resistance in children and teenagers using functional metagenomics [22]; however, in that instance they examined antibiotic resistance across a wide age range (1 month to 19 years of age), while our study provides a comprehensive insight into the infant resistome specifically at 6 months of age, demonstrating the early development of this resistome even without treatment with antibiotics. Additionally, we have demonstrated the benefits of supplementing functional screening with PCR and MIC assays, which enables the complexity of the infant resistome to be fully appreciated.

The approach applied in our study has both strengths and limitations. The strengths lie in the combination of approaches employed (functional metagenomics, PCR and MICs) to provide insights into the infant resistome. Whilst functional metagenomics enabled the detection of genes from apparently more novel sources

(including uncultured soil bacteria), PCR analysis using uncloned metagenomic DNA as a template, detected a number of gene(s) from additional sources not captured by our functional analysis. Moreover, MIC analysis highlighted the wide range of increased resistance to aminoglycosides and β -lactams conferred by each of the clone inserts. The limitations are that even using a combined metagenomic approach, we were unable to detect novel aminoglycoside or β -lactam resistance genes. It would be useful, therefore, to construct an additional infant metagenomic library using a Gram-positive host, as well as in *E. coli*, to investigate whether different resistance genes could be captured and expressed. Furthermore, it would be interesting to determine expression levels of antibiotic resistance genes harboured by the infant gut microbiota. We hope to apply such approaches in future related studies.

The findings from our study are important for several reasons. Firstly, as the infant gut is populated through complex bacterial acquisitions from maternal, dietary and environmental sources, there is the potential to control the populations colonizing infants, i.e. to maximize colonization with those bacteria that may be beneficial and pose less risk of contributing to the antibiotic resistome. Additionally, in cases where antibiotic administration is required, the route of antibiotic administration should be considered carefully when different options are available. Indeed, research using a murine model has demonstrated that antibiotic administration through intravenous injection resulted in significantly less, or the delayed development of, antibiotic resistant genes in gut microbiota populations compared to oral administration [50]. Although outside the scope of the present study, it would also seem that further investigations with respect to links between maternal antibiotic exposure and subsequent antibiotic resistant populations in the infant would be beneficial. In one previous instance, tetracycline resistance (tet^R) in the gut microbiota of one mother-infant pair was studied [21]. More specifically, resistance genes from the gut microbiota of a vaginally delivered, exclusively breastfed male infant, one month after birth, were compared with those within the gut microbiota of his mother.

References

- Sommer MOA, Dantas G, Church GM (2009) Functional characterization of the antibiotic resistance reservoir in the human microflora. *Science* 325: 1128–1131.
- Rolain J-M (2013) Food and human gut as reservoirs of transferable antibiotic resistance encoding genes. *Frontiers in Microbiology* 4: 173 doi: 10.3389/fmicb.2013.00173.
- Penders J, Stobberingh EE, Savelkoul PH, Wolfs PF (2013) The human microbiome as a reservoir of antimicrobial resistance. *Frontiers in microbiology* 4: doi: 10.3389/fmicb.2013.00087.
- Zhang L, Kinkelaar D, Huang Y, Li Y, Li X, et al. (2011) Acquired antibiotic resistance: are we born with it? *Applied and environmental microbiology* 77: 7134–7141.
- Koenig JE, Spor A, Scalfone N, Fricker AD, Stombaugh J, et al. (2011) Succession of microbial consortia in the developing infant gut microbiome. *Proceedings of the National Academy of Sciences* 108: 4578–4585.
- Fouhy F, Ross RP, Fitzgerald G, Stanton C, Cotter PD (2012) Composition of the early intestinal microbiota: Knowledge, knowledge gaps and the use of high-throughput sequencing to address these gaps. *Gut microbes* 3: 203–220.
- Marques TM, Wall R, Ross RP, Fitzgerald GF, Ryan CA, et al. (2010) Programming infant gut microbiota: influence of dietary and environmental factors. *Current opinion in biotechnology* 21: 149–156.
- Dominguez-Bello MG, Costello EK, Contreras M, Magris M, Hidalgo G, et al. (2010) Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proceedings of the National Academy of Sciences* 107: 11971–11976.
- Westerbeek EAM, van den Berg A, Laféber HN, Knol J, Fetter WPF, et al. (2006) The intestinal bacterial colonisation in preterm infants: a review of the literature. *Clinical Nutrition* 25: 361–368.
- Le Huërrou-Luron I, Biat S, Boudry G (2010) Breast-v. formula-feeding: impacts on the digestive tract and immediate and long-term health effects. *Nutr Res Rev* 23: 23–36.
- Boehm G, Moro G (2008) Structural and functional aspects of prebiotics used in infant nutrition. *Journal of Nutrition* 138: 1818S–1828S.
- Tanaka S, Kobayashi T, Songjinda P, Tateyama A, Tsubouchi M, et al. (2009) Influence of antibiotic exposure in the early postnatal period on the development of intestinal microbiota. *FEMS Immunology & Medical Microbiology* 56: 80–87.
- Fouhy F, Guinane CM, Hussey S, Wall R, Ryan CA, et al. (2012) High-throughput sequencing reveals the incomplete, short-term, recovery of the infant gut microbiota following parenteral antibiotic treatment with ampicillin and gentamycin. *Antimicrobial agents and chemotherapy* 56: 5811–5820.
- Qin X, Zerr DM, Weissman SJ, Englund JA, Denno DM, et al. (2009) Prevalence and Mechanisms of Broad-Spectrum β -Lactam Resistance in Enterobacteriaceae: a Children's Hospital Experience. *Antimicrobial agents and chemotherapy* 53: 3909–3914.
- Millar M, Philpott A, Wilks M, Whiley A, Warwick S, et al. (2008) Colonization and persistence of antibiotic-resistant Enterobacteriaceae strains in infants nursed in two neonatal intensive care units in East London, United Kingdom. *Journal of clinical microbiology* 46: 560–567.
- Gueimonde M, Salminen S, Isolauri E (2006) Presence of specific antibiotic (tet) resistance genes in infant faecal microbiota. *FEMS Immunology & Medical Microbiology* 48: 21–25.
- Kirtzalidou E, Pramatefaki P, Kotsou M, Kyriacou A (2011) Screening for lactobacilli with probiotic properties in the infant gut microbiota. *Anaerobe* 17: 440–443.
- Roy S, Viswanathan R, Singh A, Das P, Basu S (2010) Gut colonization by multidrug-resistant and carbapenem-resistant acinetobacter baumannii in neonates. *European journal of clinical microbiology & infectious diseases* 29: 1495–1500.
- Eckburg PB, Bik EM, Bernstein CN, Purdom E, Dethlefsen L, et al. (2005) Diversity of the human intestinal microbial flora. *Science* 308: 1635–1638.

Among the findings was an observation of some identical resistance genes in both populations, suggesting that the transfer of antibiotic resistant populations from the mother to the infant may occur, potentially through maternal microbiota transmission during delivery, physical contact or through breastfeeding [21]. Others have also demonstrated shared antibiotic resistance gene pools between mother and infant faecal samples [4], which may be associated with resistance genes detected in breast milk. Thus, it is likely that infant gut microbiota acquired during delivery and breastfeeding can contribute to the development of the infant resistome.

In our study, the gut microbiomes of 6 month-old infants, who had not previously been treated with antibiotics, harbored diverse antibiotic resistance genes. Metagenomic fosmid bank creation allowed functional screening for resistance genes in the gut microbiota of infants and revealed genes from diverse sources including those previously linked with uncultured bacteria. Combining this sophisticated approach with PCR techniques, provided a particularly unique insight into the presence of resistance genes in the infant gut microbiota. The results not only act as a cautionary insight into the prevalence of antibiotic resistance in infants, even in cases where no antibiotic selective pressure occurred, but also demonstrate the power of combining several metagenomic approaches to provide important insights into the infant gut resistome.

Acknowledgments

The authors wish to thank Kiera Murphy and Cian Hill for their assistance in the collection of infant faecal samples and Marc McCarthy for assistance with the QPix2-XT robotic system.

Author Contributions

Conceived and designed the experiments: FF PDC. Performed the experiments: FF. Analyzed the data: FF PDC. Contributed reagents/materials/analysis tools: FF CS PDC RPR. Wrote the paper: FF LAO BVJ RPR ACR EMD GFF CS PDC.

20. Cheng G, Hu Y, Yin Y, Yang X, Xiang C, et al. (2012) Functional screening of antibiotic resistance genes from human gut microbiota reveals a novel gene fusion. *FEMS microbiology letters* 336: 11–16.
21. de Vries LE, Vallès Y, Ageroso Y, Vaishampayan PA, García-Montaner A, et al. (2011) The gut as reservoir of antibiotic resistance: microbial diversity of tetracycline resistance in mother and infant. *PLoS ONE* 6: e21644 doi: 21610.21371/journal.pone.0021644.
22. Moore AM, Patel S, Forsberg KJ, Wang B, Bentley G, et al. (2013) Pediatric Fecal Microbiota Harbor Diverse and Novel Antibiotic Resistance Genes. *PLoS one* 8: e78822 doi:78810.71371/journal.pone.0078822.
23. Palmer C, Bik EM, DiGiulio DB, Relman DA, Brown PO (2007) Development of the human infant intestinal microbiota. *PLoS Biology* 5: e177. doi:110.1371/journal.pbio. 0050177.
24. Jones BV, Marchesi JR (2006) Transposon-aided capture (TRACA) of plasmids resident in the human gut mobile metagenome. *Nature Methods* 4: 55–61.
25. Ogilvie LA, Bowler LD, Caplin J, Dedi C, Diston D, et al. (2013) Genome signature-based dissection of human gut metagenomes to extract subliminal viral sequences. *Nature communications* 4: doi: 10.1038/ncomms3420.
26. Jones BV, Begley M, Hill C, Gahan CG, Marchesi JR (2008) Functional and comparative metagenomic analysis of bile salt hydrolase activity in the human gut microbiome. *Proceedings of the National Academy of Sciences* 105: 13580–13585.
27. Fouhy F, Ross RP, Fitzgerald GF, Stanton C, Cotter PD (2014) A degenerate PCR-based strategy as a means of identifying homologues of aminoglycoside and beta-lactam resistance genes in the gut microbiota. *BMC Microbiology* 14: 25 doi:10.1186/1471-2180-1114-1125.
28. Heuer H, Krögerrecklenfort E, Wellington E, Egan S, van Elsland J, et al. (2002) Gentamicin resistance genes in environmental bacteria: prevalence and transfer. *FEMS microbiology ecology* 42: 289–302.
29. de Fátima Silva Lopes M, Ribeiro T, Abrantes M, Figueiredo Marques JJ, Tenreiro R, et al. (2005) Antimicrobial resistance profiles of dairy and clinical isolates and type strains of enterococci. *International journal of food microbiology* 103: 191–198.
30. Schmitz F-J, Fluit AC, Gondolf M, Beyrau R, Lindenlauf E, et al. (1999) The prevalence of aminoglycoside resistance and corresponding resistance genes in clinical isolates of staphylococci from 19 European hospitals. *Journal of antimicrobial chemotherapy* 43: 253–259.
31. Bailey JK, Pinyon JL, Anantham S, Hall RM (2011) Distribution of the blaTEM gene and blaTEM-containing transposons in commensal *Escherichia coli*. *Journal of antimicrobial chemotherapy* 66: 745–751.
32. Tenover FC, Huang MB, Rasheed JK, Persing DH (1994) Development of PCR assays to detect ampicillin resistance genes in cerebrospinal fluid samples containing *Haemophilus influenzae*. *Journal of clinical microbiology* 32: 2729–2737.
33. Briñas L, Zarazaga M, Sáenz Y, Ruiz-Larrea F, Torres C (2002) β -Lactamases in ampicillin-resistant *Escherichia coli* isolates from foods, humans, and healthy animals. *Antimicrobial agents and chemotherapy* 46: 3156–3163.
34. Monstein H-J, Tärnberg M, Nilsson LE (2009) Molecular identification of CTX-M and blaOXY/K1 β -lactamase genes in Enterobacteriaceae by sequencing of universal M13-sequence tagged PCR-amplicons. *BMC Infectious Diseases* 9: 7–15.
35. Andrews JM (2001) Determination of minimum inhibitory concentrations. *Journal of antimicrobial chemotherapy* 48: 5–16.
36. Bonnet R (2004) Growing group of extended-spectrum β -lactamases: the CTX-M enzymes. *Antimicrobial agents and chemotherapy* 48: 1–14.
37. Coque TM, Oliver A, Pérez-Díaz JC, Baquero F, Cantón R (2002) Genes Encoding TEM-4, SHV-2, and CTX-M-10 Extended-Spectrum β -Lactamases Are Carried by Multiple *Klebsiella pneumoniae* Clones in a Single Hospital (Madrid, 1989 to 2000). *Antimicrobial agents and chemotherapy* 46: 500–510.
38. Morosini MI, Canton R, Martínez-Beltrán J, Negri MC, Pérez-Díaz JC, et al. (1995) New extended-spectrum TEM-type beta-lactamase from *Salmonella enterica* subsp. *enterica* isolated in a nosocomial outbreak. *Antimicrobial agents and chemotherapy* 39: 458–461.
39. Dale J, Godwin D, Mossakowska D, Stephenson P, Wall S (1985) Sequence of the OXA2 β -lactamase: comparison with other penicillin-reactive enzymes. *FEBS letters* 191: 39–44.
40. Vakulenko SB, Donabedian SM, Voskresenskiy AM, Zervos MJ, Lerner SA, et al. (2003) Multiplex PCR for detection of aminoglycoside resistance genes in enterococci. *Antimicrobial agents and chemotherapy* 47: 1423–1426.
41. Vanhoof R, Godard C, Content J, Nyssen H, Hannecart-Pokorni E (1994) Detection by polymerase chain reaction of genes encoding aminoglycoside-modifying enzymes in methicillin-resistant *Staphylococcus aureus* isolates of epidemic phage types. *Journal of medical microbiology* 41: 282–290.
42. Pallecchi L, Bartoloni A, Paradisi F, Rossolini GM (2008) Antibiotic resistance in the absence of antimicrobial use: mechanisms and implications. *Expert review of anti-infective therapy* 6: 725–732.
43. Pallecchi L, Lucchetti C, Bartoloni A, Bartalesi F, Mantella A, et al. (2007) Population structure and resistance genes in antibiotic-resistant bacteria from a remote community with minimal antibiotic exposure. *Antimicrobial agents and chemotherapy* 51: 1179–1184.
44. Bartoloni A, Pallecchi L, Rodríguez H, Fernandez C, Mantella A, et al. (2009) Antibiotic resistance in a very remote Amazonas community. *International journal of antimicrobial agents* 33: 125–129.
45. Donato JJ, Moe LA, Converse BJ, Smart KD, Berklein FC, et al. (2010) Metagenomic analysis of apple orchard soil reveals antibiotic resistance genes encoding predicted bifunctional proteins. *Applied and environmental microbiology* 76: 4396–4401.
46. Gloux K, Leclerc M, Iliozier H, L'Haridon R, Manichanh C, et al. (2007) Development of high-throughput phenotyping of metagenomic clones from the human gut microbiome for modulation of eukaryotic cell growth. *Applied and environmental microbiology* 73: 3734–3737.
47. Lorenz P, Eck J (2005) Metagenomics and industrial applications. *Nature Reviews Microbiology* 3: 510–516.
48. Rivera AM, Boucher HW (2011) Current Concepts in Antimicrobial Therapy Against Select Gram-Positive Organisms: Methicillin-Resistant *Staphylococcus aureus*, Penicillin-Resistant Pneumococci, and Vancomycin-Resistant Enterococci. Elsevier. pp.1230–1243.
49. Martiny AC, Martiny JB, Weihe C, Field A, Ellis JC (2011) Functional metagenomics reveals previously unrecognized diversity of antibiotic resistance genes in gulls. *Frontiers in microbiology* 2: 238. doi: 210.3389/fmicb.2011.00238.
50. Zhang L, Huang Y, Zhou Y, Buckley T, Wang HH (2013) Antibiotic Administration Routes Significantly Influence the Levels of Antibiotic Resistance in Gut Microbiota. *Antimicrobial agents and chemotherapy* 57: 3659–3666.