

1 Title: Early hominins in north-west Europe: a punctuated long chronology?
2 Authors & Affiliations: Rob Hosfield¹ and James Cole²
3 1. Department of Archaeology, School of Archaeology, Geography & Environmental Science,
4 University of Reading, Whiteknights, Reading, RG6 6AB, United Kingdom. E-mail:
5 r.hosfield@reading.ac.uk. ORCID ID: 0000-0001-6357-2805
6 2. School of Environment and Technology, University of Brighton, Cockcroft Building, Lewes Road,
7 Brighton, BN2 4GJ, United Kingdom. E-mail: j.n.cole@brighton.ac.uk. ORCID ID: 0000-0002-7009-
8 5303
9 Corresponding author: Rob Hosfield
10 Corresponding author address: Department of Archaeology, School of Archaeology, Geography &
11 Environmental Science, University of Reading, Whiteknights, Reading, RG6 6AB, United Kingdom.
12 Telephone: +44 (0)118 378 7632; e-mail: r.hosfield@reading.ac.uk
13 Keywords: Lower Palaeolithic; Middle Pleistocene; Europe; punctuated long chronology; life history;
14 behavioural plasticity; palaeoenvironment

15 **Abstract**

16 In light of changing views regarding the identity and evolutionary positions of Europe's Lower
17 Palaeolithic hominins, a re-consideration of the hominin occupation of North-West Europe from c. 1
18 million years ago (mya) to c. 400 thousand years ago (kya) is timely. A change in the scale and
19 character of the overall European Palaeolithic record around c. 800-600 kya has been well
20 documented and argued over since the mid-1990s. Hominin expansion into the European north-
21 west, potentially from southern Europe, Africa or south-western Asia, has been linked to the
22 introduction of a new lithic technology in the form of the biface. We evaluate three potential drivers
23 for this northern range expansion: changing palaeo-climatic conditions, the emergence of an
24 essentially modern human life history, and greater hominin behavioural plasticity. Our evaluation
25 suggests no major changes in these three factors during the c. 800-600 kya period other than
26 enhanced behavioural plasticity suggested by the appearance of the biface. We offer here a model
27 of hominin occupation for north-west Europe termed the 'punctuated long chronology' and suggest
28 that the major changes in the European Lower Palaeolithic record that occur at a species wide level
29 may post-date, rather than precede, the Anglian Glaciation (marine isotope stage (MIS) 12).

30

31 **Highlights**

- 32 • European Lower Palaeolithic includes key shift in hominin distribution.
- 33 • Shift from below 45° north to broader range including higher latitudes.
- 34 • Changes in palaeoenvironmental conditions or hominin life history not key factors.
- 35 • Increasingly plastic behaviour may have enabled hominin range expansion.

36

37 **1. Introduction**

38 The hominin occupation of North-West Europe is a complex and important story for understanding
39 hominin abilities to adapt to novel environments and the implementation of the plastic behaviours
40 that were first developed on the savannahs of Africa and Eurasia (Dennell, 2003; Dennell and

41 Roebroeks, 2005; Grove, 2011; Potts, 2013). North-West Europe is important due to the unique
42 challenges that faced hominins within the higher latitude belt in terms of that region's distinctive
43 flora, fauna and seasonality. The earliest hominin occupation of Europe has traditionally been seen
44 through the long, short, and modified short chronologies (Carbonell et al., 1996; Dennell and
45 Roebroeks, 1996; Dennell, 2003; Roebroeks and van Kolfschoten, 1994; Table 1), recently enriched
46 through the Eurasian perspective (Dennell, 2004).

Model	Long Chronology (Carbonell et al., 1996)	Short Chronology (Roebroeks and van Kolfschoten, 1994)		Modified Short Chronology (Dennell and Roebroeks, 1996)
Summary	Europe occupied in the Early Pleistocene, and in some cases before c. 1.5 mya	"No undisputable proof for human occupation of Europe prior to about 500,000 years ago" (Roebroeks and van Kolfschoten, 1994: 500)		"Hominids might have occasionally moved into southern Europe well before 500,000 years ago, as and when conditions permitted" (Dennell and Roebroeks, 1996: 540)
Evidence				
Timeframe	Before 780,000 years ago (pan-Europe)	Before 500,000 years ago (pan-Europe)	After 500,000 years ago (pan-Europe)	Before 780,000 years ago (Southern Europe)
Archaeological signature	Small assemblages (typically 10s and 100s of artefacts)	Small series of isolated pieces selected from a natural pebble background	Large collections from excavated knapping floors with conjoinable material	Small assemblages (typically 10s and 100s of artefacts)
Context	Cave(?) and open-air sites (pan-Europe)	Disturbed secondary contexts	Primary context sites (fine-grained matrix)	Cave(?) and open-air sites (Iberia)
Assemblage character	Flake and pebble tools	Contested 'primitive' assemblages	Uncontested Acheulean and non-Acheulean	Flake and pebble tools

			industries	
Human Remains	Occasional	No human remains	Human remains common	Occasional

47

48 **Table 1:** Summarising the long, short and modified short chronologies, as originally proposed (Carbonell et al., 1996; Dennell and Roebroeks, 1996; Gamble,

49 1999; Roebroeks and van Kolfschoten, 1994).

50 The long and short chronologies have been hampered by the resolution and scale of the
51 archaeological record and uncertainty whether evidence for hominin presence and absence in
52 North-West Europe prior to 800 kya is a genuine behavioural pattern or the result of research and/or
53 taphonomic bias (Roebroeks and van Kolfschoten, 1994). However, discoveries at a number of sites
54 including Dmanisi, Georgia (Lordkipanidze et al., 2007); Orce and Atapuerca, Spain (Carbonell et al.,
55 2005; Carbonell et al., 1995; Toro-Moyano et al., 2013); Pirro Nord, Italy (Arzarello et al., 2007);
56 [Pont-de-Lavaud and Lunery-Rosières, France](#) (Despriée et al., 2011); and Happisburgh III, UK (Ashton
57 et al., 2014; Parfitt et al., 2010) have been instrumental in challenging the notion of a hominin
58 absence in Europe pre-800 kya. Happisburgh III has also challenged the notion of early hominin
59 intolerance for harsh climatic conditions. There are however, still clear arguments to be made in
60 regards to the sustainable nature of early forays [substantially](#) above the 45°N mark (MacDonald et
61 al., 2012).

62

63 A related question concerns the increasingly complex role of *Homo antecessor* in the earliest human
64 occupation of north-west Europe. It is *Homo antecessor* that has been put forward as one of the
65 contenders for the pre-800 kya human presence at Happisburgh III (Ashton et al., 2014). Although
66 their fossil evidence is limited to the Iberian Peninsula, it is possible that a series of short-lived
67 dispersal events by this species above 45°N may have been the cause of the sparse archaeological
68 record in Europe before 800 kya.

69

70 The pace and rhythms of early hominin dispersal in Europe are therefore varied and challenging to
71 decipher. What is clear however is that hominin development must have occurred in conjunction
72 with changes in climate and environment. It is the understanding of the specific climatic and
73 environmental challenges to hominin survival in north-west Europe, and potential solutions to those
74 challenges, which forms the focus of this paper.

75

76 2. How to survive Pleistocene North West Europe

77 Delays in the occupation of northern Eurasia, above c. 45°N, after the initial dispersals or range
78 expansions of *H. erectus* (*s. lato*) beyond Africa, suggest that a sustained presence in the north
79 presented significant new challenges to Lower Palaeolithic hominins. Modern ethnographic,
80 ecological and climatological studies would suggest that those challenges were principally increased
81 seasonality, lower mean temperatures, contrasting oceanic and continental zones, reduced plant
82 food resources, and more dispersed and clustered animal food resources (Gamble, 1986; Hosfield,
83 2016; Kelly, 1995; Roebroeks, 2001, 2006). How might these challenges be met? The 'solutions' can
84 be grouped into two broad categories: firstly, increased animal food contributions to the diet, with
85 implications for the reliability of foraging strategies, the sizes of territories and the knowledge
86 required of landscapes and resources, hominin mobility, social group organisation, and technology.
87 Secondly, enhanced insulation, which can be cultural (encompassing some or all of pyrotechnology,
88 whether managed or opportunistic, 'clothing' or other forms of body covering, and shelter) and/or
89 physiological (e.g. elevated BMR [basal metabolic rate], increased muscle mass, body hairs, elevated
90 brown adipose tissue levels; but *cf.* Aiello and Wheeler, 1995; Aiello and Wheeler, 2003).

91
92 These 'solutions' can be most easily described as changing hominin behaviours (i.e. increasingly
93 efficient dietary and insulation strategies). Yet our clearest material insights into hominin behaviour
94 (the lithic record) shows relatively little evidence for major changes across the duration of the Lower
95 Palaeolithic (Klein, 2000), and while Acheulean technology may be absent from the repertoire of the
96 earliest Europeans, it already had a long history in Africa and western Asia by the time of those first
97 dispersals into the Mediterranean. We therefore propose three other potential forms of change
98 which may also be key factors in the pre-/post-800kya settlement patterns in northern Eurasia
99 (Table 2), and ~~will~~ explore these within a European framework: (i) changes in the palaeoclimate and
100 palaeoenvironmental context; (ii) changes in hominin life history; and (iii) changes in behavioural
101 plasticity.

Period	Chronology	Sites
Later Early Pleistocene	c. 1.3–0.78mya	Happisburgh III, Lunery-Rosières, Pont-de-Lavaud
Early Middle Pleistocene	c. 0.78–0.5mya	Abbeville, Boxgrove, Happisburgh I, High Lodge, Kärlich G, La Genetière, La Noira, Miesenheim, Pakefield, Warren Hill
Later Middle Pleistocene	c. 0.5–0.3mya	Beeches Pit, Bilzingsleben, Clacton, Hoxne, Schöningen, Soucy, Swanscombe,

102

103 **Table 2:** Key chronological phases and selected northern European sites (after Ashton, 2015).

104

105 **2.1 The palaeoenvironmental context**

106 To what extent may changing palaeoclimatic and palaeoenvironmental conditions underpin the
107 changes in scale in the northern European settlement record? Such arguments have been widely
108 explored. Turner (1992) emphasised the changing large carnivore guild around 500kya and the
109 gradual emergence of the 'mammoth steppe' after MIS 12 (Guthrie, 1990), linked to the changing
110 length of glacial/interglacial cycles during the Mid-Pleistocene Transition (Head and Gibbard, 2005).
111 However, many of these models have tended to primarily emphasise long-term trends. While these
112 are undoubtedly important, the emphasis in this paper is upon the daily, seasonal and yearly
113 challenges to hominins: e.g. low winter temperatures, the degree of seasonality, short-term climatic
114 shifts (e.g. on decadal scales), the diversity or homogeneity of habitats at a local scale, and the
115 condition and availability of plant and animal foods across the seasons. These are increasingly well
116 understood through micro-fauna (e.g. Coope, 2006a), palynology, and the multi-disciplinary
117 investigations of key sites (Bigga et al., 2015; García and Arsuaga, 2011; Parfitt et al., 2010; Parfitt et
118 al., 2005; Rodríguez et al., 2011; Urban and Bigga, 2015), enabling us to consider the nature of the
119 hominin life experience 'on the ground'. We are taking this approach not least because while the
120 MIS record emphasises the increasing stability, length, marked oscillations and decreasing

121 temperature of European glacial/interglacial cycles during the Middle Pleistocene, intra-site records
122 reveal that stability is less apparent at the sub-stage scale (Ashton et al., 2008a; Urban and Bigga,
123 2015) and may also have been less apparent at the scale of a hominin lifetime too. The key question
124 is can we see clear evidence for a shift in hominin environments across the c. 800-600kya boundary
125 that might favour a more sustained northern presence?

126

127 **2.2 Changes in hominin life history**

128 Changing life history may be a second key factor. The current European fossil record suggests an
129 apparent species contrast pre- and post-800-600kya, with *H. antecessor* (Carbonell et al., 1995)
130 replaced by *H. heidelbergensis* (Stringer, 2012) across the chronological division. It may well also be
131 significant that the Sima de los Huesos fossils, dating to c. 430kya, are now argued to be early
132 Neanderthals, or closely related to the latter's ancestors (Meyer et al., 2016b). Why might life
133 history be significant? The modern human life history stages (childhood, juvenile, adolescence,
134 adulthood) are associated with the unusual collection of traits which characterise humans:
135 prolonged gestation, growth and maturation, extremely short inter-birth intervals, helpless
136 newborns, a short period of breastfeeding/early weaning, extended offspring dependency, an
137 adolescence growth-spurt, delayed reproduction, and the menopause (Bogin and Smith, 1996;
138 Schwartz, 2012).

139

140 Of particular importance to the occupation of the north may be the emphasis in the human model of
141 early weaning: this strategy places infants at risk, as they are unskilled at finding appropriate foods
142 (of sufficient high quality to fuel brain growth, but also suitable for small, deciduous teeth), are
143 essentially defenceless, and can be competing with other adults (Aiello and Key, 2002; Kennedy,
144 2003). It therefore has notable dietary strategy implications, but it also has significant implications
145 for infant care. Since early weaning is associated with shorter inter-birth intervals, other forms of
146 childcare are required for the 'weanlings', in order to avoid excessive DEE (daily energy expenditure)


147 loads on the large-bodied and large-brained *Homo* females (i.e. normal DEE + gestation/lactation +
148 nursing of 'weanlings'; Aiello and Key, 2002). This alloparenting can come from grandmothers (e.g.
149 Hawkes et al., 1998) or older siblings, or in the form of increased male provisioning.

150

151 As noted above, a change to a fully modern life history would therefore bring costs, but the addition
152 of an adolescence stage can also bring specific opportunities to improve selective fitness. In a
153 broader context, extended growth development would enable socialisation – learning the 'rules' of
154 social life (e.g. hierarchies of food access, recognising the moods and emotions of individuals), made
155 complicated both by larger communities, as predicted by neocortex size and the social brain
156 hypothesis (e.g. Gamble et al., 2014), and perhaps also by periods of individuals' absences, arising
157 from dispersed resource distributions and social fission-fusion in the seasonal mid- and high-
158 latitudes (Couzin and Laidre, 2009). Kennedy (2003) has argued that adolescence would enable
159 youngsters who were close to maturity to observe and participate in the social, sexual and infant-
160 care practices that are critical to success in adulthood. Bogin and Smith (1996) have similarly
161 emphasised adolescence as a period when parenting skills could be practiced, perhaps resulting in
162 the greater survival of their own offspring later in life. MacDonald (2007) has stressed the knowledge
163 demands of a hunting life, from animal behaviour and their signs, to animal-animal and animal-plant
164 relationships, to the topography, vegetation and seasonality of the environment. Foraging for plant
165 resources, while stationary, would also require a diverse set of ecological knowledge. Adolescence
166 would offer an opportunity to develop much of this base-line knowledge, and the associated skills –
167 not least because the nature and content of learning will vary with age, as both the mind and the
168 body develops (MacDonald, 2007). Thus a modern human model of life history might produce sub-
169 adults and adults with the skills and knowledge to meet the social, ecological and technological
170 challenges of the north.

171

172 We can discuss life history and its evolution amongst extinct hominins (Fig. 1) because of the
 173 evidence for strong correlations between a range of life history traits: brain size, body size, age at
 174 sexual maturity, age at first birth, gestational length, lifespan, and dental development (Kennedy,
 175 2003). The critical question from a European Lower Palaeolithic perspective would therefore seem
 176 to be: can we detect any evidence for adolescence in particular, and a truly human model of growth
 177 and development in the hominins of the late Early Pleistocene and early Middle Pleistocene?
 178



179
 180 **Fig. 1:** Inferred life history stages for selected hominin species (Bogin, 1999: fig. 4.9). *H. erect*⁻¹ =
 181 early *H. erectus* (adult brain sizes of 850-900cc); *H. erect*⁻² = late ~~*H. erectus*~~ *H. erectus* (adult brain sizes
 182 upto 1100cc); *A. afar* = *A. afarensis*; *A. Africa* = *A. africanus*.

183
 184 **2.3 Changes in behavioural plasticity**

185 In addition to hominin genotypic adaptations to the north-west European palaeoclimatic and
 186 palaeoenvironmental conditions (although some traditionally considered “cold-adapted” hominin

187 traits have recently been questioned (e.g. Rae et al., 2011) hominin range expansion must also have
188 been characterised by phenotypic adaptations to external factors. Archaeologically speaking it is the
189 phenotypic adaptations of behaviour that are the most visible indicators of behavioural plasticity.


190

191 It has been suggested for some time that organisms that are both mobile and long-lived are less
192 likely to adapt to local conditions through the selection of genetically based traits (Potts, 1998, 2013;
193 van Schaik, 2013). In effect, the suggestion is therefore that when such organisms encounter
194 variable climatic conditions they are more likely to adapt to those conditions at a phenotypic level
195 rather than a genotypic one (van Schaik, 2013). However, it would also appear that the plastic
196 behavioural strategy may fail if local environments change too rapidly (Van Buskirk, 2012; van
197 Schaik, 2013). This in turn suggests that behavioural plasticity can only carry a species so far in its
198 effectiveness in negotiating novel or rapidly changing environments.

199

200 Another important aspect to consider in regards to behavioural plasticity are the links to cognitive
201 capacity and brain size (van Schaik, 2013). From a primate perspective behavioural plasticity derives
202 from learning through mental mechanisms linked to varying degrees of cognitive complexity
203 (Shettleworth, 2010). These processes have been tightly linked to brain size and therefore suggest
204 that behavioural plasticity should be included in the costs of supporting larger brains (van Schaik,
205 2013). Therefore, it follows that the degree of behavioural plasticity present within an organism may
206 correspondingly be limited by overall brain size. Within the hominin fossil record it has been
207 suggested previously that there are recognisable step changes in hominin brain size (Maslin et al.,
208 2014; McNabb and Cole, 2015; Shultz et al., 2012; Fig. 2) at c. 1.8mya, 1mya, 400–200kya, and
209 100kya, which punctuate periods of stasis in hominin brain expansion. Shultz et al. (2012) further
210 highlight that the increase in hominin brain size c. 400–200 kya may be as a result of migrating
211 hominin species into Eurasia as the step changes are not mirrored in Africa. However, from Figure 2

212 it could be cautiously suggested (based on the limited sample available) that there may also be an
 213 increase in hominin brain size within *Homo heidelbergensis* c. 800–600 kya.
 214



215
 216 **Fig. 2: Showing Hominin cranial volumes (cm³) at over the last 2myr a (A) and 1mya (B) scale. The**
 217 **fossil specimens and species are divided by location** (data after Berger et al., 2015; Brown et al.,
 218 2004; Shultz et al., 2012; Thackeray, 2015).

219
 220 Indeed, if behavioural plasticity and brain size are linked then there is a possibility that a more
 221 sustained hominin presence in north-west Europe may require a threshold level of both plasticity
 222 and brain size. Therefore, examining the fossil and behavioural signatures of Pleistocene hominin
 223 activity in north-west Europe against the context of changing environmental conditions should
 224 provide some insight as to the degree of plasticity present, and whether an increase in behavioural

225 plasticity coinciding with an increase in brain size is significant enough of an advantage to lead to a
226 more sustained hominin presence after c. 600kya.

227

228 3. Discussion

229 Following the three broad themes presented above, we will now examine and review the evidence
230 for each.

231

232 3.1 Palaeoenvironment

233 Kahlke et al. (2011) argue that 1.2–0.9mya (the late Early Pleistocene) was a period of less uniform
234 climate cycles, varying in both duration and intensity. These climate cycles would have been
235 associated with new mammal communities, increasing varieties of habitats, large riverine systems,
236 and relatively mild and humid conditions, especially in the Mediterranean and the west. As Kahlke et
237 al. (2011) have previously argued, these conditions seemingly favoured the expanding occupation of
238 the Mediterranean zone at sites such as Barranco León and Fuentenueva-3 (Orce), Sima del Elefante
239 (TE9), Pont-de-Lavaud, Lunery-Rosières, Le Vallonet III (although the character of the lithic
240 assemblage has been strongly critiqued; Roebroeks and van Kolfschoten, 1994) and Monte Poggiolo
241 (Leroy et al., 2011), while dispersal into the north was restricted to favourable interglacials of low
242 seasonality and high habitat diversity, for example at Happisburgh III (Parfitt et al., 2010). By
243 contrast, the period after 0.9mya was increasingly controlled by 100kyr periodicity, with longer and
244 relatively stable climatic intervals, progressive temperature decline and increasing aridity, marked
245 oscillations between highly contrasting landscapes, greater seasonality, and increasingly specialized
246 mammal communities (Kahlke et al., 2011: 1389). These changes ~~become~~became especially marked
247 after c. 500kya (Berger and Loutre, 1991; Lisiecki and Raymo, 2005). Perhaps critically from the
248 perspective of hominin distribution and dispersal, these changes were accompanied by regional
249 shifts in characteristic habitat types (Table 3).

Region	Prevailing habitats (inferred from large mammal communities)				
Apennine Peninsula	High variety of open / forested habitats	Open woodland / steppe			
Iberian Peninsula	Open savannah / lightly forested habitats	Open woodland, tree savannah / steppe			
Western Europe	High variety of open / forested habitats	Woodland / steppe	Variety of open to forested habitats		
Northwest Europe	Woodland	Woodland with open patches / mixed habitats	Steppe / woodland		
Central Europe	High variety of open / forested habitats	Steppe / woodland with open patches	Steppe-tundra / woodland		
Eastern Europe	High variety of forest steppe / open steppe habitats	Open steppe / forest steppe	Steppe-tundra / woodland		
Chronology (mya)	1.2	1.0	0.8	0.6	0.4

250

251 **Table 3:** Prevailing habitat characteristics in Europe over the last 1.2mya (re-drawn after Kahlke et
252 al., 2011: Fig. 6).

253

254 What might these regional differences have meant in terms of hominin life on the ground? Can we
255 see any evidence of significant change over time? Offering a northern Iberian perspective from
256 Atapuerca, Rodríguez et al. (Rodríguez et al., 2011) have argued for an absence of harsh conditions

257 across the long time-span of the Sima del Elefante, Gran Dolina, and Galería sites. This was based on
258 a wide range of evidence: small vertebrates ([amphibians, squamates and small mammals](#)), large
259 mammals, and vegetation ([Table 4](#)). Rodríguez et al. (2011) conclude that these prevalent temperate
260 conditions may be associated with a continuous early hominin presence prior to c. 0.5mya, and
261 perhaps reflect a *H. antecessor* preference for warmer, less variable conditions (see also Agustí et al.,
262 2009).

263

264 So what changes in terms of hominin habitats? The answer from south-western Europe, using
265 Atapuerca as a case study, may be not much. Rodríguez et al.'s (2011) admittedly tentative
266 environmental reconstructions (Table 4) suggest little change between the Early and Middle
267 Pleistocene, while Garcia & Arsuaga's (2011) analysis of the Sima de los Huesos evidence suggests a
268 savannah-like open woodland, dating to c. 430kya on the basis of the most recent dating (Arnold et
269 al., 2014). But what is the view from the north?

<u>Evidence</u>	<u>Key Patterns (Early & Middle Pleistocene)</u>	
<u>Small vertebrates</u>	<ul style="list-style-type: none"> • <u>Significant woodland component in herpetofauna</u> • <u>Small mammals suggest moist, open habitats</u> 	
<u>Large mammals</u>	<ul style="list-style-type: none"> • <u>All of temperate affinity</u> • <u>None unequivocally indicating cold or arid environments</u> 	
<u>Vegetation</u>	<ul style="list-style-type: none"> • <u>Continuous presence of Mediterranean taxa & mesic trees</u> • <u>One possible cold steppe landscape phase</u> 	
<u>Palaeotemperature</u> <u>(amphibian & squamate assemblages)</u>	<ul style="list-style-type: none"> • <u>Always warmer than Burgos today</u> • <u>“Cold”/“warm” period differences of only c. 2°C</u> 	
<u>Annual precipitation</u>	<ul style="list-style-type: none"> • <u>750mm (dry)-1049mm (humid), compared to c. 570mm in present</u> 	
<u>Environment types</u>	<u>Early Pleistocene</u> <ul style="list-style-type: none"> • <u>Mediterranean open woodland</u> • <u>Open woodland and steppe</u> • <u>Open arid (?)</u> • <u>Open woodland and moist meadows</u> • <u>Temperate open woodlands and meadows with lagoons</u> 	<u>Middle Pleistocene</u> <ul style="list-style-type: none"> • <u>Humid temperate open woodland</u> • <u>Mediterranean open woodland</u> • <u>Moist meadows with scattered trees</u> • <u>Open woodland and meadows</u> • <u>Steppe</u>

270

271 **Table 4:** Inferred environments at Atapuerca in the Early and Middle Pleistocene (after Rodríguez et
272 al., 2011: table 9).

273

274 There is evidence of a tolerance of cooler conditions as a whole (e.g. Hosfield, 2011: table 6), gained
275 from palaeotemperature estimates and environmental descriptors from a range of key northern

276 sites with evidence of hominin activity. These offer a valuable perspective on possible changes in
 277 hominin tolerances and/or preferences in northern Europe across the c. 800-600kya 'boundary'
 278 (Table 5). Summer and winter palaeo-temperatures reveal no major changes across the boundary as
 279 a whole, although there is perhaps some evidence for slightly harsher conditions in the MIS 13/15
 280 sites, in contrast to the MIS 11 sites and Pakefield. This might be significant, and slightly counter-
 281 intuitive, as MIS 13 sees a marked increase in the number and size of sites across northern Europe
 282 (e.g. examples in Tables 5 & 6, plus Warren Hill, Highlands Farm, Valdoe, Kent's Cavern, Abbeville,
 283 Miesenheim, and Kärlich G; see also Ashton, 2015).
 284

Site	T _{min} (°C)	T _{max} (°C)	Evidence ¹	Age (MIS)	Source
Early Pleistocene					
Happisburgh III (Bed E)	-3 - 0	+16 - \pm 18	Coleoptera	Late 25 or late 21	(Ashton and Lewis, 2012; Parfitt et al., 2010)
Early Middle Pleistocene					
Pakefield (Bed Cii-Ciii)	-6 - +4	+17 - \pm 23	Coleoptera	17 or later 19	(Ashton and Lewis, 2012; Coope, 2006b)
Boxgrove (Unit 4c & Freshwater Silt Bed ≈ Units 4b & 4c)	-4 - +4	+15 - \pm 20	Ostracods (MOTR) & Herpetofauna (MCR)	13	(Ashton and Lewis, 2012; Holman, 1999; Holmes et al., 2010)

Happisburgh I (Organic Mud)	-11 - -3	+12 - <u>+15</u>	Coleoptera	13?	(Ashton and Lewis, 2012; Coope, 2006b)
High Lodge (Bed C1)	-4 - +1	+15 - <u>+16</u>	Coleoptera	13?	(Coope, 2006b)
Waverley Wood (Channel 2, Organic Mud)	-	+10 - <u>+15</u>	Coleoptera	13 or 15	(Coope, 2006b; Shotton et al., 1993)
Brooksby (Redland's Brooksby Channel)	-10 - +2	+15 - <u>+16</u>	Coleoptera	13 or 15	(Coope, 2006b)
Later Middle Pleistocene					
Barnham (Unit 5c; Holl)	-	+17 - <u>+18</u>	Herpetofauna	11c	(Holman, 1998)
Hoxne (Stratum D ⁵ ; Holl ⁶)	-10 - +6	+15 - <u>+19</u>	Coleoptera	11c	(Ashton et al., 2008a; Coope, 1993)
Bilzingsleben II	-0.5 - +3	+20 - <u>+25</u>	Mollusca & ostracods ²	11	(Mania, 1995; Mania and Mania, 2003)
Schöningen 13 II-4	-4 - -1	+16	Mollusca & ostracods	9	(Urban and Bigga, 2015)
Present Day Comparisons					

East Anglia (present day) ³	-0.7 - +6.9	+14.2 - <u>+</u> 18.0	-	-	-
Bilzingsleben (present day) ⁴	<u>-3.3 - +2.0</u>	+ <u>12.4 -</u> <u>+22.7</u>	-	-	-

285

286

Table 5: Winter and summer temperature estimates for Early and Middle Pleistocene British and

287

German sites (with selected present day equivalents for reference). ¹Sensitivity tests on coleoptera-

288

based MCR procedures suggest that winter temperature estimates are usually too warm (Pettitt and

289

White, 2012: 35). ²The specific source of the palaeo-temperature estimates is not stated, but the

290

fauna includes molluscs and ostracods; ³East Anglian data based on Met Office annual mean

291

seasonal temperatures (1910–2016;

292

http://www.metoffice.gov.uk/pub/data/weather/uk/climate/datasets/Tmean/date/East_Anglia.txt);

293

⁴Bilzingsleben data [based on Deutscher Wetterdienst \(German Weather Service\) monthly mean](#)

294

[January and July temperatures \(1951–2017; Erfurt-Weimar station;](#)

295

<https://www.dwd.de/DE/leistungen/klimadatendeutschland/klarchivtagmonat.html>)

296

297

Site	Palaeoenvironmental description	Age (MIS)	Source
Early Pleistocene			
Pont-de-Lavaud	Closed deciduous forest; warm, wet climate	30	(Messenger et al., 2011)
Happisburgh III (Unit C)	Cool temperate; heathland & coniferous forest	Late 25 or late 21	(Ashton and Lewis, 2012)
Happisburgh III (Units D & E)	Cool temperate; dominated by coniferous forest	Late 25 or late 21	(Ashton and Lewis, 2012)

Early Middle Pleistocene			
Pakefield (Units Aiv-Ciii)	Mediterranean-type climate; marshy ground with reed-beds & alder carr, surrounded by deciduous forest	17 or later 19	(Ashton and Lewis, 2012)
Happisburgh I (Organic Mud-Organic Silt Sand)	Cool climate; marshy edge of slow-flowing river, with surrounding coniferous forest	13?	(Ashton and Lewis, 2012)
High Lodge (Units B2-C2)	Cool-temperate climate; marshy ponds, heath & grassland on river floodplain, surrounded by pine & spruce forest	13?	(Ashton and Lewis, 2012)
Boxgrove (Unit 4b)	Coastal mudflats, with occasional drying out & spread of grassland	13	(Ashton and Lewis, 2012)
Boxgrove (Unit 4c & Q1B pond deposits)	Temperate climate, with cooler winters than present; grassland, scrub & mixed woodland, with spring-fed pools	13	(Ashton and Lewis, 2012)
Boxgrove (Unit 8)	Interstadial environment	Later 13-early 12	(Ashton and Lewis, 2012)
Boxgrove (Unit 11)	Periglacial, with ameliorations in climate	Later 13-early 12	(Ashton and Lewis, 2012)
Later Middle Pleistocene			
Bilzingsleben II	Fully temperate, with subcontinental influences; mixed oak woodland & shrub associations, alternating with steppe meadows	11	(Mania and Mania, 2003)
Barnham (Unit 5c;	Temperate climate; mix of grassland &	11c	(Ashton and

Holl)	deciduous woodland on edge of slow-flowing river		(Lewis, 2012)
Beeches Pit (Unit 4; Holll)	Fully temperate environment; spring-fed pools with deciduous woodland	11c	(Ashton and Lewis, 2012)
Beeches Pit (Unit 3b; Hollb-c)	Temperate climate; shallow to stagnant pools with calcareous grassland and surrounding open woodland	11c	(Ashton and Lewis, 2012)
Hoxne (Unit B1)	Cooler temperatures (<i>cf.</i> units D & E); probable birch & pine forest; some grassland areas	11a	(Ashton and Lewis, 2012)
Swanscombe (Lower Loam; Hollb-c)	Grassy floodplain with shallow pools & occasional flooding; mixed oak forest nearby, with hazel & alder	11c	(Ashton and Lewis, 2012)
Schöningen 13 II-4	Mosaic swamp, with reed-like and sedge vegetation; regional context: highly continental dry steppe/open forest; tree taxa dominated by pine & birch	9	(Stahlschmidt et al., 2015; Urban and Bigga, 2015)


298

299 **Table 6:** Palaeoenvironmental reconstructions for Early and Middle Pleistocene British sites and
300 German sites.

301

302 The general trend after MIS 13 is towards full or cool-temperate environments, mixing deciduous or
303 coniferous woodland with open grass and/or heathland, and water sources (Table 6). As Bigga et al.
304 (2015) have noted at Schöningen, such environments provide a diverse range of potential plant
305 foods, meeting both nutritional (e.g. vitamin C in pine and birch bark, berries of common bearberry,
306 European elder, raspberry, and leaves of *Ranunculus* and *Chenopodium*) and seasonal needs (Figure

307 3). Evidence of animal butchery also suggests few changes in preferred prey across the c. 800-600kya
 308 boundary, or in comparison between northern and southern Europe (Table 7).
 309



310
 311 **Figure 3:** Seasonal variations in selected potential plant food sources identified at Schöningen 13 II-4
 312 (Bigga et al., 2015).
 313

Period	Species	Butchery Evidence	Sites
Early Pleistocene	<i>Bison</i> sp.	Defleshing; disarticulation;	Gran Dolina TD6 (Saladié et al., 2011)
	<i>B. menneri</i>	dismemberment; evisceration;	
	<i>B. voigtstedtensis</i>	filleting; skinning	Sima del Elefante (Huguet et al., 2013)
	<i>C. elaphus</i>		
	<i>C. s.l. nestii vallonnetensis</i>		

	<i>Cervid sp.</i> <i>Cercopithecidae</i> <i>Eucladoceros</i> <i>Equus sp.</i> <i>Hippopotamus</i> <i>S. etruscus</i> <i>S. hundsheimensis</i> <i>U. dolinensis</i> <i>V. praeglacialis</i>		
Early Middle Pleistocene	<i>Bos</i> or <i>Bison</i> sp. <i>C. capreolus</i> <i>C. elaphus</i> <i>E. ferus</i> <u><i>S. hundsheimensis</i></u> <i>U. deningeri</i>	Cut-marks; defleshing; dismemberment; filleting; marrow bone breakage; skinning	Boxgrove (Bello et al., 2009; Parfitt and Roberts, 1999) Happisburgh I (Ashton et al., 2008b) Westbury (Andrews and Ghaleb, 1999)
Later Middle Pleistocene	<i>Bos</i> or <i>Bison</i> sp. <i>C. elaphus</i> <i>E. ferus</i>	Boning; cut-marks; defleshing; dismemberment; filleting; marrow bone breakage; skinning	Barnham (Parfitt, 1998) Schöningen (Van Kolfschoten et al., 2015a; Voormolen, 2008) Hoxne (Stopp, 1993)

314

315 **Table 7:** European sites with butchery evidence.

316

317 What is perhaps more noticeable are the conditions at the Early Pleistocene sites of Happisburgh III
318 and Pont-de-Lavaud, and at the MIS 17/19 site of Pakefield. The latter two sites both support a
319 model of temporary northern excursions in line with favourable conditions. As other authors have
320 previously noted (Parfitt et al., 2010), Happisburgh III stands out in the pre-800-600kya record, albeit
321 within a very small sample of early northern sites.

322

323 Overall, combining general palaeo-environmental trends and the conditions at hominin sites
324 suggests that the post-800-600kya expansions in the occupation of the north occur at an intuitively
325 odd time - before the favourable MIS 11 interglacial but after the milder conditions of the later Early
326 Pleistocene.

327

328 **3.2 Life history**

329 Turning away from environmental conditions, can we detect any evidence for childhood and
330 adolescence, and a truly human model of growth and development, in the European Lower
331 Palaeolithic? Of critical importance in beginning to resolve this question for Europe has been the *H.*
332 *heidelbergensis* and *H. antecessor* fossil material from the Sima de los Huesos and Gran Dolina
333 localities at Atapuerca (Bermúdez de Castro et al., 1997; Carbonell et al., 1995; these analyses pre-
334 dated the recent re-attribution of the Sima de los Huesos material as possible early Neanderthals).
335 Initial analysis of the dental material from TD-6 hominins 1 & 3 and Sima hominin 18 suggested that
336 both European species are similar to modern humans in their dental development, implying both
337 prolonged maturation and new life history stages (childhood and adolescence; Bermúdez de Castro
338 et al., 2003), although *H. antecessor* is not as derived in some respects (e.g. delayed M3 calcification;
339 Bermúdez de Castro et al., 1999). By extension that would also suggest relatively short birth
340 intervals, high rates of postnatal brain growth, extended offspring dependency, marked adolescent
341 growth spurt, and delayed reproductive cycles. Expanded analysis of enamel formation rates for
342 various species of *Homo* also suggested prolonged dental growth in *H. antecessor* and *H.*

343 *heidelbergensis* (Ramirez Rozzi and Bermudez de Castro, 2004). Although their mean crown
344 formation times were concluded to be shorter than in Upper Palaeolithic–Mesolithic *H. sapiens*
345 samples, they were slower than those of Neanderthals, whose crown formation was estimated to be
346 15% quicker than modern humans. The data also suggested only slightly slower growth rates in *H.*
347 *heidelbergensis* (n=106; 21 individuals) compared to *H. antecessor* (n=8; 4 individuals; *ibid*: fig. 1).

348

349 However, these interpretations have been challenged to some extent by Robson and Wood (2008),
350 who have argued that *H. antecessor*'s crown formation rates are not modern human-like (although
351 the tooth eruption timings are). They also noted that the crown formation rates of *H.*

352 *heidelbergensis* are more akin to *H. antecessor* than modern humans, a view supported by Rozzi &
353 Bermudez de Castro's data (2004: fig. 1). Contrasts with *H. sapiens* are also evident in Neanderthals'
354 more rapid dental development patterns (Ramirez Rozzi and Bermudez de Castro, 2004; Smith et al.,
355 2007), although suggested differences between those two species' overall rates of growth have
356 recently been challenged (Rosas et al., 2017). Overall Robson and Wood (2008) conclude that life
357 history shifts are likely to have been piece-meal (with body mass shifts pre-dating dental
358 developments; see also Krovitz et al., 2003), and also that dental development rates and timings do
359 not always correlate well with other life history variables.

360

361 These various models and data suggest that European Lower Palaeolithic hominins, whether *H.*
362 *antecessor* or *H. heidelbergensis*, are likely to have adopted at least a human-like model of
363 reproduction, growth and development, although childhood may not have been as prolonged as in
364 *H. sapiens*. However, it is worth considering whether life history may have become more human-like
365 with *H. heidelbergensis* (i.e. more prolonged maturation and more extended childhood and
366 adolescence stages), given the overlaps in endocranial volume between *H. sapiens*, *H.*
367 *neanderthalensis* and *H. heidelbergensis*, but not *H. antecessor* (Robson and Wood, 2008: fig. 6), and
368 the delayed M3 calcification and long I2 and C crown formation times in *H. antecessor* (Bermúdez de

369 Castro et al., 2003; Bermúdez de Castro et al., 1999). As noted above the human life history pattern
370 offers the opportunity to learn skills and experiences that may have been critical in the seasonal
371 latitudes of Europe (e.g. the extended practice argued to be required by prepared platform
372 techniques; Stout et al., 2014). But differences between *H. heidelbergensis* and *H. antecessor* are
373 likely to have been relatively small and it is not immediately obvious as to how and why small
374 extensions in the childhood and adolescence stages, and other associated life-history changes,
375 would facilitate the post-600kya expansion in northern Europe. In many ways, the suggested
376 appearance of an adolescence stage in *H. antecessor*, and perhaps also *H. erectus* (Krovitz et al.,
377 2003; but cf. Bogin and Smith, 1996, and Fig. 1) looks more significant, and may well relate to the
378 earlier, first excursions into southern Europe.

379

380 **3.3 Behavioural plasticity**

381 Within the Lower Palaeolithic record, the biface represents a shift to a multi-functional tool that is
382 easily transportable and adaptable (through resharpening and reworking) to a variety of situations
383 (Moncel et al., 2015). There are a range of additional behavioural and cognitive changes that biface
384 manufacture represents, including indications of greater planning depth, not only in tool production,
385 use and curation, but also a more liberated attitude to landscape use and resource acquisition that
386 potentially underpinned an expansion in group size and more complex group dynamics (Gowlett et
387 al., 2012). Much of these additional bonuses may come from the acquisition of a sustained ability for
388 abstract thought at a species level (Cole, 2015a; Cole, 2015b) contributing to a greater degree of
389 sustained behavioural plasticity as brain sizes increase (Fig. 2). Therefore, tracking the earliest
390 appearance of biface sites in north-west Europe could provide a useful insight into hominin
391 behavioural changes. We follow Ashton (2015) in associating the bifaces of north-west with *Homo*
392 *heidelbergensis*, as demonstrated at sites such as Boxgrove and Arago (Moncel et al., 2015).

393

394 Moncel et al. (2015) have produced an excellent review of biface manufacture in north-west Europe
395 from c. 700–600kya which suggests a more extensive biface presence in north-west Europe from
396 500kya (Table 8).

Age (MIS)	Contexts	Raw Materials	Characteristic Biface Morphology	Additional Flake Tools	Sites
17-13	Diamicton; fluvial gravel; fluvial sand; fluvial silts; lagoonal silts; palaeosol	Andesite; siliceous; flint; quartzite	Pointed (incl. crude forms); ovate; biface- cleaver; cordiforms	Denticulates; notches; scrapers	Boxgrove (Q1B); Brandon Fields; Happisburgh I; High Lodge (Bed E; Sands & Gravels); La Noira; Maidscross Hill; Warren Hill; Waverley Wood
12-10	Gravel; slope deposit; fluvial sand; fluvial gravel; tufa; pool; colluvial; alluvial	Flint	Pointed (incl. crude forms); backed biface; cleaver; cordiform; ovate; sub-triangular	Denticulates; endscrapers; notches; scrapers; pebble tools; retouched flakes	Cagny-La-Garenne I (Cxb, Cxv, Lg, Lj & Ca assemblages); Cagny-La-Garenne II (13, 14 & J assemblages); Rue De Cagny (Series 3); La Celle; Barnham; Elveden; Beeches Pit; Swanscombe (LMG & UMG); Hoxne (Upper & Lower Industries); Ferme de l'Épinette (MS assemblage); St- Pierre-lès-Elbeuf

397

398 **Table 8:** Summary table of early north-western biface sites, context, raw material, biface form and additional flake tool types (data from Moncel et al.,
399 2015).

400

401 When looking at the early biface assemblages of north-west Europe (Table 8) it can be seen that
402 there is a diversity of biface form within the assemblages and the sites tend to cluster around fluvial
403 locations, often at the convergence of ecological zones (Table 6). This is perhaps not surprising as
404 this pattern of hominin site location has been repeated throughout the Old World, probably
405 reflecting the preference of hominins in exploiting a range of resources in the vein of the generalist
406 (Grove, 2011; Potts, 2013). What is interesting is a pattern noticed by Moncel et al. (2015), that the
407 early biface sites in north-west Europe tend to have a relatively low ratio of bifaces to other tools
408 when compared to comparable sites in the Levant or Africa (although there are a few notable
409 exceptions, e.g. Boxgrove). While there are taphonomic complications to this pattern (e.g. variable
410 collecting histories), two potentially key trends have been highlighted (Moncel et al., 2015):

411

412 Firstly, the Early Middle Pleistocene sites (MIS 17-13) tend to have not only a range of biface forms
413 made from both core and flake blanks, but they are also characterised by a range of raw material
414 exploitations depending on local availability. This may indicate that the Early Middle Pleistocene
415 biface makers were not locked into a single raw material but flexible enough in cognition and
416 experience to work a range of lithologies when necessary. Alternatively, of course, it may equally
417 suggest a limited ability for planning depth with a more immediate focus on utilising suitable raw
418 materials as they occur in the local landscape. In addition, raw material seems to have differing
419 impacts on the degree of shaping and size of bifaces present between southern British and northern
420 French sites which may go some way to explaining the difference in artefact morphology. However,
421 a common occurrence between these sites are the dual compositions of biface assemblages
422 consisting of thick, crudely pointed bifaces with irregular cutting edges and more comprehensively
423 worked bifaces with regular cutting edges and invasive removals.

424

425 Secondly, the late Middle Pleistocene (MIS 12-10) assemblages in contrast seem to be characterised
426 by higher proportions of thinner bifaces with fine retouched edges, although overall shaping is still

427 varied within assemblages. Across the Table 8 sites large flakes could have been produced for blank
428 production, but it would appear that few were. This may in turn reflect the suitability and increasing
429 visibility of good flint sources in the landscape as well as reflect the hominins' ability to read and
430 interpret the landscape around them. Another possible change during the late Middle Pleistocene in
431 Europe (in contrast to the early Middle Pleistocene) is that the biface sites may be supported in the
432 toolkit by a larger range of flake tool types (Table 8), although it is acknowledged that collection
433 biases and varying typological schemes also complicate this interpretation.

434

435 Certainly, the patterns of hominin behaviour seen in the early biface sites of north-west Europe are
436 diverse, but they tend to be explained in terms of the differences in local raw material and overall
437 site function even if there are broad similarities in the shaping processes in biface manufacture
438 (Moncel et al., 2015). We would suggest that these differences and similarities in biface shaping are
439 probably more a consequence of local technological and functional decisions rather than cultural
440 factors at this time.

441

442 Nonetheless the European biface package would also seem to suggest a range of additional
443 technological developments – although it is not the position here that they are European
444 innovations or that the biface per se enabled northern dispersals. Rather, it is the overall suggested
445 behavioural package that may have facilitated the sustainable tipping point for hominin dispersal
446 into north-west Europe. Preservation records are difficult for non-lithic technologies but there are
447 preserved wooden spears from Clacton c. 400 kya (Warren, 1911) and spears and bone tools from
448 Schöningen c. 300–337 kya (Richter and Krbetschek, 2015; Van Kolfschoten et al., 2015b). While
449 recent palaeogenetic evidence raises the possibility that these artefacts may be the preserve of a
450 later speciation event (Meyer et al., 2016a), the mosaic character of Middle Pleistocene hominin
451 evolution combined with the Boxgrove puncture wound evidence supports a position that the early

452 biface makers of north-west Europe may have had an extensive lithic and non-lithic tool kit including
453 spears.

454

455 In addition, hints of hide processing can be seen in the elaborate scrapers from High Lodge, Warren
456 Hill, Brandon Fields and Maidscross Hill (Moncel et al., 2015) which would presumably indicate a
457 possible use for them as clothing or shelter components. However, direct evidence for such
458 behaviour is sadly lacking. It is also clear that the later biface-making hominins had the controlled
459 use of fire from well-known sites such as Beeches Pit, Ménez Drégan and possibly Aroeira at c. 400
460 kya (Daura et al., 2017; Gowlett, 2006; Molines et al., 2005; Preece et al., 2006). Although some
461 argue that these are isolated instances of controlled fire use in Europe (Roebroeks and Villa, 2011)
462 they are supported by dates for controlled fire use in other parts of Old World since c. 800 kya at
463 least (Goren-Inbar et al., 2004).

464

465 The biface itself has often been seen to represent something more than just a functional tool with
466 explanations varying from butchery, mechanisms of sexual selection and aesthetic markers to
467 cultural mediators and objects used to negotiate the landscapes and social-scapes of the Acheulean
468 world (Gamble, 1998; Hopkinson and White, 2005; Kohn and Mithen, 1999; Machin et al., 2007;
469 Machin and Mithen, 2004; McPherron, 2000; Porr, 2005; Saragusti et al., 1998; Wenban-Smith,
470 2004; White, 1998; Wynn, 1995). We agree with aspects of the above and emphasise again that it is
471 not the biface per se that allowed hominins to have a more sustained presence in north-west
472 Europe, but perhaps the full behavioural package that accompanied biface manufacture.

473

474 Why then did it take so long for biface makers to enter north-west Europe when there is clearly
475 evidence for these artefacts stretching back to 1.7 million years ago in Africa (Diez-Martin et al.,
476 2015) and first appearing in south-western Asia soon after (Pappu et al., 2011)? We suggest here
477 that not only do you need the biface package (with all the social and technological benefits that

478 includes) but you also need a threshold level of behavioural plasticity that is tied to a brain size of
479 over 1100cm³ at a sustained species level for hominin populations to persistently survive the
480 difficulties of the European north-west. Since Moncel et al. (2015) and Table 8 demonstrate that
481 bifaces are present in Europe from at least c. 700 kya, we suggest that this plasticity package must
482 have been present across the early/late Middle Pleistocene boundary (MIS 13/12).

483

484 **4. Conclusion**

485 Our interpretation of the palaeoenvironmental (life on the ground) and palaeoanthropological (life
486 history) records from north-western Europe across the c. 800-600kya period has struggled to find a
487 clear pattern of changing conditions before and after this 'boundary'. By contrast, the archaeological
488 evidence does highlight the appearance of the biface by c. 700kya, although it does not become
489 widespread until MIS 13. The status of this possible behavioural boundary is newly noteworthy in
490 light of the recent palaeo-genetic studies that have pushed back the date of the last common
491 ancestor between modern humans and Neanderthals to c. 700 kya (Meyer et al., 2016a). This makes
492 the apparent step-change in the European archaeological record ever more intriguing. Perhaps such
493 changes may simply relate to the appearance of a new species (*H. heidelbergensis?*) able to cope
494 with the rigours of the north-west, although it remains uncertain as to whether such a species is an
495 *in situ* development or a dispersal into Europe from elsewhere.

496

497 However, there may be a subtler and more nuanced pattern to the archaeological record, reflecting
498 a multi-phased hominin occupation of north-west Europe, which we dub here the 'punctuated long
499 chronology' (Table 9), which builds on the modified short chronology (Dennell and Roebroeks, 1996).
500 The first phase, contemporary with the pre-1 million year flake and core occupations of the southern
501 European zone exemplified by Dmanisi, Orce and Atapuerca, is characterised by northern absence.
502 Shortly after 1 million years ago there would appear to be small scale, fragmented dispersals of
503 hominins further north into cooler climates such as at Happisburgh III, equipped with a flake and

504 core lithic tool kit. This is succeeded by the appearance of biface-producing hominins between c.
505 700-500kya, possibly originating from regions outside Europe. These biface-makers had a varied
506 lithic and organic tool kit, incorporating the behavioural plasticity package within a broad geographic
507 range, but would still seem to be limited in terms of overall population numbers. This in turn may
508 reflect the continuing challenges of a hominin population adapting to the unique conditions of
509 Pleistocene north-west Europe. We suggest that this second phase of small-scale, biface-making
510 populations, although important, has tended to be over-emphasised in interpretations of the north-
511 west European Palaeolithic record due to the quality of preservation at occasional key sites such as
512 Boxgrove. In contrast, we suggest here that the major behavioural change (the plasticity package)
513 occurs at a species level, leading to demographic growth, during a third phase after MIS 12. This can
514 tentatively be supported by the changing richness, and diversity in tool kits, of the artefactual record
515 (e.g. the terrace archives of the Thames, Solent and Somme; Ashton and Hosfield, 2010; Ashton and
516 Lewis, 2002; Tuffreau and Antoine, 1995) and evidence for cooler climate tolerances (e.g. Hoxne and
517 Cagny-la-Garenne). Perhaps most tantalisingly, such a shift might also map onto the latest
518 palaeogenetic timings for the origins of the early Neanderthals at around c. 430kya (Meyer et al.,
519 2016a). Further evidence is certainly need to test this hypothesis, and we suggest an enhanced
520 understanding of the quantities of artefacts at different periods, normalised with reference to
521 sample biases and site taphonomy (e.g. Ashton and Hosfield, 2010; Ashton and Lewis, 2002), as one
522 possible approach.

Phase	1	2	3	4
Summary	Only southern Europe occupied.	Continued occupation of southern Europe, with small-scale, fragmented dispersals into northern Europe (e.g. Happisburgh III).	Continued occupation of southern and northern Europe. Emerging behavioural plasticity package (possibly originating from outside Europe), but limited overall population numbers, especially in the north.	Full behavioural plasticity package and expanded population numbers across Europe.
Timeframe	Before 1mya	c. 1mya-700kya	c. 700-500kya	c. 500-300kya
Archaeological signature	Small assemblages (typically 10s and 100s of artefacts)	Small assemblages (typically 10s and 100s of artefacts)	Larger individual assemblages (100s and 1000s of artefacts)	Large assemblages (1000s and 10000s of artefacts), with multiple large sites
Context	Cave(?) and open-air sites	Cave(?) and open-air sites	Cave and open-air sites	Cave and open-air sites
Assemblage character	Flake and pebble tools, with potential organic tools	Flake and pebble tools, with potential organic tools	Bifaces and flake tools (limited range), with demonstrated	Bifaces and flake tools (expanded range), with demonstrated

			organic tools	organic tools
Human Remains	Very occasional	Occasional	Occasional	Occasional, but with abundant remains at individual sites

523

524 **Table 9:** A punctuated long chronology for the hominin occupation of Europe.

525

526 **Acknowledgements:**

527 The ideas in this paper benefitted greatly from the AHRC-funded Network meeting “Coping with
528 climate: the legacy of *Homo heidelbergensis*”, paper presentations at the Royal Anthropological
529 Institute conference “*Anthropology, Weather and Climate Change*” and the *Palaeolithic-Mesolithic*
530 2016 meeting at the British Museum: we thank all those who attended and commented on our
531 ideas. We are also grateful for the thoughtful comments provided by the two anonymous reviewers.
532 All remaining ideas and errors are our own.

533

534 **Funding:**

535 This work was supported by the Arts & Humanities Research Council (grant number:
536 AH/N007514/1).

537

538 **References:**

- 539 Agustí, J., Blain, H.A., Cuenca-Bescós, G., Bailon, S., 2009. Climate forcing of first hominid
540 dispersal in Western Europe. *Journal of Human Evolution* 57, 815-821.
- 541 Aiello, L.C., Key, C., 2002. Energetic consequences of being a *Homo erectus* female.
542 *American Journal of Human Biology* 14, 551-565.
- 543 Aiello, L.C., Wheeler, P., 1995. The expensive tissue hypothesis: the brain and the digestive
544 system in human and primate evolution. *Current Anthropology* 36, 199 - 221.
- 545 Aiello, L.C., Wheeler, P., 2003. Neanderthal Thermoregulation and the Glacial Climate, in:
546 van Andel, T.H., Davies, W. (Eds.), *Neanderthals and modern humans in the European*
547 *landscape during the last glaciation*. McDonald Institute for Archaeological Research,
548 Cambridge, pp. 147-166.
- 549 Andrews, P., Ghaleb, B., 1999. Taphonomy of the Westbury Cave bone assemblages, in:
550 Andrews, P., Cook, J., Carrant, A., Stringer, C.B. (Eds.), *Westbury Cave: The Natural*
551 *History Museum Excavations, 1976–1984*. Western Academic & Specialist Press Limited,
552 Bristol, pp. 87–126.
- 553 Arnold, L.J., Demuro, M., Parés, J.M., Arsuaga, J.L., Aranburu, A., Bermúdez de Castro,
554 J.M., Carbonell, E., 2014. Luminescence dating and palaeomagnetic age constraint on
555 hominins from Sima de los Huesos, Atapuerca, Spain. *Journal of Human Evolution* 67, 85-
556 107.
- 557 Arzarello, M., Marcolini, F., Pavia, G., Pavia, M., Petronio, C., Petrucci, M., Rook, L.,
558 Sardella, R., 2007. Evidence of earliest human occurrence in Europe: the site of Pirro Nord
559 (Southern Italy). *Naturwissenschaften* 94, 107-112.
- 560 Ashton, N., Hosfield, R., 2010. Mapping the Human Record in the British Early Palaeolithic:
561 Evidence from the Solent River System. *Journal of Quaternary Science* 25, 737-753.

562 Ashton, N., Lewis, S.G., 2002. Deserted Britain: Declining Populations in the British Late
563 Middle Pleistocene. *Antiquity* 76, 388-396.

564 Ashton, N., Lewis, S.G., 2012. The environmental contexts of early human occupation of
565 northwest Europe: The British Lower Palaeolithic record. *Quaternary International* 271, 50-
566 64.

567 Ashton, N., Lewis, S.G., De Groote, I., Duffy, S.M., Bates, M.R., Bates, R., Hoare, P., Lewis,
568 M., Parfitt, S.A., Peglar, S., Williams, C., Stringer, C., 2014. Hominin Footprints from Early
569 Pleistocene Deposits at Happisburgh, UK. *PLoS One* 9, e88329.

570 Ashton, N., Lewis, S.G., Parfitt, S.A., Penkman, K.E., Coope, G.R., 2008a. New evidence for
571 complex climate change in MIS 11 from Hoxne, Suffolk, UK. *Quaternary Science Reviews*
572 27, 652-668.

573 Ashton, N.M., 2015. Ecological niches, technological developments and physical adaptations
574 of early humans in Europe: the handaxe-*heidelbergensis* hypothesis, in: Coward, F., Hosfield,
575 R.T., Pope, M., Wenban-Smith, F.F. (Eds.), *Settlement, Society and Cognition in Human*
576 *Evolution: Landscapes in Mind*. Cambridge University Press, Cambridge, pp. 138-153.

577 Ashton, N.M., Parfitt, S.A., Lewis, S.G., Coope, G.R., Larkin, N., 2008b. Happisburgh Site 1
578 (TG388307), in: Candy, I., Lee, J.R., Harrison, A.M. (Eds.), *The Quaternary of Northern East*
579 *Anglia Field Guide*. Quaternary Research Association, London, pp. 151–156.

580 Bello, S.M., Parfitt, S.A., Stringer, C.B., 2009. Quantitative micromorphological analyses of
581 cut marks produced by ancient and modern handaxes. *Journal of Archaeological Science* 36,
582 1869-1880.

583 Berger, A., Loutre, M.F., 1991. Insolation values for the climate of the last 10 million years.
584 *Quaternary Science Reviews* 10, 297-317.

585 Berger, L.R., Hawks, J., de Ruiter, D.J., Churchill, S.E., Schmid, P., Delezene, L.K., Kivell,
586 T.L., Garvin, H.M., Williams, S.A., DeSilva, J.M., Skinner, M.M., Musiba, C.M., Cameron,
587 N., Holliday, T.W., Harcourt-Smith, W., Ackermann, R.R., Bastir, M., Bogin, B., Bolter, D.,
588 Brophy, J., Cofran, Z.D., Congdon, K.A., Deane, A.S., Dembo, M., Drapeau, M., Elliott,
589 M.C., Feuerriegel, E.M., Garcia-Martinez, D., Green, D.J., Gurtov, A., Irish, J.D., Kruger, A.,
590 Laird, E.W., Marchi, D., Meyer, M.R., Nalla, S., Negash, E.W., Orr, C.M., Radovčić, D.,
591 Schroeder, L., Scott, J.E., Throckmorton, Z., Tocheri, M.W., VanSickle, C., Walker, C.S.,
592 Wei, P., Zipfel, B., 2015. *Homo naledi*, a new species of the genus *Homo* from the Dinaledi
593 Chamber, South Africa. *eLIFE*, 4:e09560.

594 Bermúdez de Castro, J.M., Arsuaga, J.L., Carbonell, E., Rosas, A., Martínez, I., Mosquera,
595 M., 1997. A hominid from the Lower Pleistocene of Atapuerca: possible ancestor to
596 Neanderthals and Modern Humans. *Science* 276, 1392–1395.

597 Bermúdez de Castro, J.M., Ramirez Rozzi, F.V., Martinon-Torres, M., Sarmiento Perez, S.,
598 Rosas, A., 2003. Patterns of dental development in Lower and Middle Pleistocene hominins
599 from Atapuerca (Spain), in: Thompson, J.L., Krovitz, G.E., Nelson, A.J. (Eds.), *Patterns of*
600 *Growth and Development in the Genus Homo*. Cambridge University Press, Cambridge, pp.
601 246-270.

602 Bermúdez de Castro, J.M., Rosas, A., Carbonell, E., Nicolás, M.E., Rodríguez, J., Arsuaga,
603 J.L., 1999. A modern human pattern of dental development in Lower Pleistocene hominids
604 from Atapuerca-TD6 (Spain). *Proceedings of the National Academy of Sciences* 96, 4210-
605 4213.

606 Bigga, G., Schoch, W.H., Urban, B., 2015. Paleoenvironment and possibilities of plant
607 exploitation in the Middle Pleistocene of Schöningen (Germany). *Insights from botanical*
608 *macro-remains and pollen*. *Journal of Human Evolution* 89, 92-104.

609 Bogin, B., 1999. *Patterns of Human Growth*, 2nd ed. Cambridge University Press,
610 Cambridge.

611 Bogin, B., Smith, B.H., 1996. Evolution of the human life cycle. *American Journal of Human*
612 *Biology* 8, 703-716.

613 Brown, P., Sutikna, T., Morwood, M.J., Soejono, R.P., Jatmiko, E., Saptomo, E.W., Due,
614 R.A., 2004. A new small-bodied hominin from the Late Pleistocene of Flores, Indonesia.
615 *Nature* 431, 1055-1061.

616 Carbonell, E., Bermúdez de Castro, J.M., Arsuaga, J.-L., Allue, E., Bastir, M., Benito, A., I,
617 C., Canals, T., Diez, J.C., van der Made, J., Mosquera, M., Ollé, A., Pérez-González, A.,
618 Rodríguez, J., Rodríguez, X.P., Rosas, A., Rosell, J., Sala, R., Vallverdú, J., Vergés, J.M.,
619 2005. An Early Pleistocene hominin mandible from Atapuerca-TD6, Spain. *Proceedings of*
620 *the National Academy of Sciences* 102, 5674-5678.

621 Carbonell, E., Bermudez de Castro, J.M., Arsuaga, J.-L., Diez, J.C., Rosas, A., Cuenca-
622 Bescos, G., Sala, R., Mosquera, M., Rodríguez, X.P., 1995. Lower Pleistocene hominids and
623 artifacts from Atapuerca-TD6 (Spain). *Science* 269, 826-830.

624 Carbonell, E., Mosquera, M., Rodríguez, X.P., Sala, R., 1996. The first human settlement of
625 Europe. *Journal of Anthropological Research* 52, 107-114.

626 Cole, J., 2015a. Examining the Presence of Symmetry within Acheulean Handaxes: A Case
627 Study in the British Palaeolithic. *Cambridge Archaeological Journal* 24, 713-732.

628 Cole, J., 2015b. Handaxe Symmetry in the Lower and Middle Palaeolithic: Implications for
629 the Acheulean Gaze in: Coward, F., Hosfield, R., Pope, M., Wenban-Smith, F. (Eds.),
630 *Settlement, Society and Cognition in Human Evolution: Landscapes in Mind*. Cambridge
631 University Press, Cambridge, pp. 234-257.

632 Coope, G.R., 1993. Late-Glacial (Anglian) and Late-Temperate (Hoxnian) Coleoptera, in:
633 Singer, R., Gladfelter, B.G., Wymer, J.J. (Eds.), *The Lower Palaeolithic Site at Hoxne,*
634 *England*. University Chicago Press, Chicago, pp. 156-162.

635 Coope, G.R., 2006a. Insect faunas associated with Palaeolithic industries from five sites of
636 pre-Anglian age in central England. *Quaternary Science Reviews* 25, 1738-1754.

637 Coope, G.R., 2006b. Insect Faunas Associated with Palaeolithic Industries from Five Sites of
638 Pre-Anglian Age in Central England. *Quaternary Science Reviews* 25, 1738-1754.

639 Couzin, I.D., Laidre, M.E., 2009. Fission–fusion populations. *Current Biology* 19, R633-
640 R635.

641 Daura, J., Sanz, M., Arsuaga, J.L., Hoffmann, D.L., Quam, R.M., Ortega, M.C., Santos, E.,
642 Gómez, S., Rubio, A., Villaescusa, L., Souto, P., Mauricio, J., Rodrigues, F., Ferreira, A.,
643 Godinho, P., Trinkaus, E., Zilhão, J., 2017. New Middle Pleistocene hominin cranium from
644 Gruta da Aroeira (Portugal). *Proceedings of the National Academy of Sciences* 114, 3397-
645 3402.

646 Dennell, R., Roebroeks, W., 1996. The earliest colonization of Europe: the short chronology
647 revisited. *Antiquity* 70, 535-542.

648 Dennell, R.W., 2003. Dispersal and colonisation, long and short chronologies: how
649 continuous is the Early Pleistocene record for hominids outside East Africa? *Journal of*
650 *Human Evolution* 45, 421-440.

651 Dennell, R.W., 2004. Hominid dispersals and Asian biogeography during the Lower and
652 Early Middle Pleistocene, ca. 2.0 - 0.5 mya. *Asian Perspectives* 43, 205-226.

653 Dennell, R.W., Roebroeks, W., 2005. Out of Africa: An Asian perspective on early human
654 dispersal from Africa. *Nature* 438, 1099-1104.

655 Despriée, J., Voinchet, P., Tissoux, H., Bahain, J.-J., Falguères, C., Courcimault, G., Dépont,
656 J., Moncel, M.-H., Robin, S., Arzarello, M., Sala, R., Marquer, L., Messager, E., Puaud, S.,
657 Abdessadok, S., 2011. Lower and Middle Pleistocene human settlements recorded in fluvial
658 deposits of the middle Loire River Basin, Centre Region, France. *Quaternary Science*
659 *Reviews* 30, 1474-1485.

660 Diez-Martin, F., Yustos, P.S., UribeArrea, D., Baquedano, E., Mark, D.F., Mabulla, A.,
661 Fraile, C., Duque, J., Perez-Gonzalez, A., Yravedra, J., Egeland, C.P., Organista, E.,
662 Dominguez-Rodrigo, M., 2015. The Origin of the Acheulean: The 1.7 Million-Year-Old Site
663 of FLK West, Olduvai Gorge (Tanzania). *Scientific Reports* 5, 17839.

664 Gamble, C., Gowlett, J., Dunbar, R.I.M., 2014. *Thinking Big: How the Evolution of Social*
665 *Life Shaped the Human Mind*. Thames & Hudson, London.

666 Gamble, C.S., 1986. *The Palaeolithic Settlement of Europe*. Cambridge University Press,
667 Cambridge.

668 Gamble, C.S., 1998. Handaxes and Palaeolithic Individuals, in: Ashton, N., Healy, F., Pettitt,
669 P. (Eds.), *Stone Age Archaeology: Essays in honour of John Wymer*. Oxbow Monograph
670 Oxford.

671 Gamble, C.S., 1999. *The Palaeolithic Societies of Europe*. Cambridge University Press,
672 Cambridge.

673 García, N., Arsuaga, J.L., 2011. The Sima de los Huesos (Burgos, northern Spain):
674 palaeoenvironment and habitats of *Homo heidelbergensis* during the Middle Pleistocene.
675 *Quaternary Science Reviews* 30, 1413-1419.

676 Goren-Inbar, N., Alperson, N., Kislev, M.E., Simchoni, O., Melamed, Y., Ben-Nun, A.,
677 Werker, E., 2004. Evidence of Hominin Control of Fire at Geshert Benot Ya'akov, Israel.
678 *Science* 304, 725-727.

679 Gowlett, J.A.J., 2006. The Early Settlement of Northern Europe: Fire History in the Context
680 of Climate Change and the Social Brain. *C. R. Palevol* 5, in press.

681 Gowlett, J.A.J., Gamble, C.S., Dunbar, R.I.M., 2012. Human Evolution and the Archaeology
682 of the Social Brain. *Current Anthropology* 53, 693-722.

683 Grove, M., 2011. Speciation, diversity, and Mode 1 technologies: The impact of variability
684 selection. *Journal of Human Evolution* 61, 306-319.

685 Guthrie, R.D., 1990. *Frozen Fauna of the Mammoth Steppe*. University of Chicago Press,
686 Chicago.

687 Hawkes, K., O'Connell, J.F., Jones, N.G.B., Alvarez, H., Charnov, E.L., 1998.
688 Grandmothering, menopause, and the evolution of human life histories. *Proceedings of the*
689 *National Academy of Sciences* 95, 1336-1339.

690 Head, M.J., Gibbard, P.L., 2005. Early-Middle Pleistocene transitions: an overview and
691 recommendation for the defining boundary. *Geological Society, London, Special Publications*
692 247, 1-18.

693 Holman, J.A., 1998. The herpetofauna. The interglacial mammalian fauna from Barnham, in:
694 Ashton, N.M., Lewis, S.G., Parfitt, S.A. (Eds.), *Excavations at the Lower Palaeolithic Site at*
695 *East Farm, Barnham, Suffolk 1989-94*. The British Museum, London, pp. 101-106.

696 Holman, J.A., 1999. Herpetofauna, in: Roberts, M.B., Parfitt, S.A. (Eds.), *Boxgrove: a*
697 *Middle Pleistocene hominid site at Earham Quarry, Boxgrove, West Sussex*. English
698 *Heritage, London*, pp. 181-187.

699 Holmes, J.A., Atkinson, T., Fiona Darbyshire, D.P., Horne, D.J., Joordens, J., Roberts, M.B.,
700 Sinka, K.J., Whittaker, J.E., 2010. Middle Pleistocene climate and hydrological environment
701 at the Boxgrove hominid site (West Sussex, UK) from ostracod records. *Quaternary Science*
702 *Reviews* 29, 1515-1527.

703 Hopkinson, T., White, M.J., 2005. The Acheulean and the handaxe: structure and agency in
704 the Palaeolithic, in: Gamble, C.S., Porr, M. (Eds.), *The Hominid Individual in Context:*
705 *Archaeological investigations of Lower and Middle Palaeolithic landscapes, locales and*
706 *artefacts*. Routledge, London and New York, pp. 13 - 28.

707 Hosfield, R., 2016. Walking in a Winter Wonderland? Strategies for Early and Middle
708 Pleistocene Survival in Midlatitude Europe. *Current Anthropology* 57, 653-682.

709 Hosfield, R.T., 2011. The British Lower Palaeolithic of the early Middle Pleistocene.
710 *Quaternary Science Reviews* 30, 1486 - 1510.

711 Huguet, R., Saladié, P., Cáceres, I., Díez, C., Rosell, J., Bennàsar, M., Blasco, R., Esteban-
712 Nadal, M., Gabucio, M.J., Rodríguez-Hidalgo, A., Carbonell, E., 2013. Successful
713 subsistence strategies of the first humans in south-western Europe. *Quaternary International*
714 295, 168-182.

715 Kahlke, R.-D., García, N., Kostopoulos, D.S., Lacomat, F., Lister, A.M., Mazza, P.P.A.,
716 Spassov, N., Titov, V.V., 2011. Western Palaeartic palaeoenvironmental conditions during
717 the Early and early Middle Pleistocene inferred from large mammal communities, and
718 implications for hominin dispersal in Europe. *Quaternary Science Reviews* 30, 1368-1395.

719 Kelly, R.L., 1995. *The Foraging Spectrum: Diversity in Hunter-Gatherer Lifeways*.
720 Smithsonian Institution Press, Washington.

721 Kennedy, G.E., 2003. Palaeolithic Grandmothers? Life History Theory and Early Homo.
722 *Journal of the Royal Anthropological Institute* 9, 549-572.

723 Klein, R.G., 2000. Archeology and the evolution of human behavior. *Evolutionary*
724 *Anthropology* 9, 17-36.

725 Kohn, M., Mithen, S.J., 1999. Handaxes: products of sexual selection? *Antiquity* 73, 518-
726 526.

727 Krovitz, G.E., Thompson, J.L., Nelson, A.J., 2003. Hominid growth and development from
728 australopithecines to Middle Pleistocene *Homo*, in: Thompson, J.L., Krovitz, G.E., Nelson,
729 A.J. (Eds.), *Patterns of Growth and Development in the Genus Homo*. Cambridge University
730 Press, Cambridge, pp. 271-292.

731 Leroy, S.A.G., Arpe, K., Mikolajewicz, U., 2011. Vegetation context and climatic limits of
732 the Early Pleistocene hominin dispersal in Europe. *Quaternary Science Reviews* 30, 1448-
733 1463.

734 Lisiecki, L.E., Raymo, M.E., 2005. A Pliocene-Pleistocene stack of 57 globally distributed
735 benthic $\delta^{18}\text{O}$ records. *Paleoceanography* 20.

736 Lordkipanidze, D., Jashashvili, T., Vekua, A., Ponce de León, M.S., Zollikofer, C.P.E.,
737 Rightmire, G.P., Pontzer, H., Ferring, R., Oms, O., Tappen, M., Bukhsianidze, M., Agusti, J.,
738 Kahlke, R., Kiladze, G., Martínez-Navarro, B., Mouskhelishvili, A., Nioradze, M., Rook, L.,
739 2007. Postcranial evidence from early Homo from Dmanisi, Georgia. *Nature* 449, 305-310.

740 MacDonald, K., 2007. Ecological hypotheses for human brain evolution: evidence for skill
741 and learning processes in the ethnographic literature on hunting, in: Roebroeks, W. (Ed.),
742 *Guts and Brains: an integrative approach to the hominin record*. Leiden University Press,
743 Leiden, pp. 107-132.

744 MacDonald, K., Martínón-Torres, M., Dennell, R.W., Bermúdez de Castro, J.M., 2012.
745 Discontinuity in the record for hominin occupation in south-western Europe: Implications for
746 occupation of the middle latitudes of Europe. *Quaternary International* 271, 84-97.

747 Machin, A., Hosfield, R.T., Mithen, S.J., 2007. Why are some handaxes symmetrical?
748 Testing the influence of handaxe morphology on butchery effectiveness. *Journal of*
749 *Archaeological Science* 34, 883 - 893.

750 Machin, A., Mithen, S., 2004. Reply to McNabb et al 2004. The large cutting tools from the
751 South African Acheulean and the questions of social traditions. *Current Anthropology* 45,
752 653 - 677.

753 Mania, D., 1995. The earliest occupation of Europe: the Elbe-Saale region (Germany), in:
754 Roebroeks, W., van Kolfschoten, T. (Eds.), *The Earliest Occupation of Europe*. Proceedings
755 of the European Science Foundation Workshop at Tautavel (France) 1993. Leiden University
756 Press & European Science Foundation, Leiden, pp. 85-102.

757 Mania, D., Mania, U., 2003. Bilzingsleben — Homo erectus, his culture and his environment.
758 The most important results of research, in: Burdukiewicz, J.M., Ronen, A. (Eds.), Lower
759 Palaeolithic Small Tools In Europe And The Levant. BAR, Oxford, pp. 29-48.

760 Maslin, M.A., Brierley, C.M., Milner, A.M., Shultz, S., Trauth, M.H., Wilson, K.E., 2014.
761 East African climate pulses and early human evolution. *Quaternary Science Reviews* 101, 1-
762 17.

763 McNabb, J., Cole, J., 2015. The mirror cracked: Symmetry and refinement in the Acheulean
764 handaxe. *Journal of Archaeological Science: Reports* 3, 100-111.

765 McPherron, S.P., 2000. Handaxes as a measure of the mental capabilities of early hominids.
766 *Journal of Archaeological Science* 27, 655 - 663.

767 Messenger, E., Lebreton, V., Marquer, L., Russo-Ermolli, E., Orain, R., Renault-Miskovsky,
768 J., Lordkipanidze, D., Despriée, J., Peretto, C., Arzarello, M., 2011. Palaeoenvironments of
769 early hominins in temperate and Mediterranean Eurasia: new palaeobotanical data from
770 Palaeolithic key-sites and synchronous natural sequences. *Quaternary Science Reviews* 30,
771 1439-1447.

772 Meyer, M., Arsuaga, J.-L., de Filippo, C., Nagel, S., Aximu-Petri, A., Nickel, B., Martinez, I.,
773 Gracia, A., Bermúdez de Castro, J.M., Carbonell, E., Viola, B., Kelso, J., Prüfer, K., Pääbo,
774 S., 2016a. Nuclear DNA sequences from the Middle Pleistocene Sima de los Huesos
775 hominins. *Nature* 531, 504-507.

776 Meyer, M., Arsuaga, J.-L., Filippo, C., Nagel, S., Aximu-Petri, A., Nickel, B., Martinez, I.,
777 Gracia, A., de Castro, J.M.B., Carbonell, E., Viola, B., Kelso, J., Prüfer, K., Pääbo, S., 2016b.
778 Nuclear DNA Sequences from the Middle Pleistocene Sima de los Huesos Hominins. *Nature*
779 531, 504-507.

780 Molines, N., Monnier, J.-L., Hinguant, S., Hallegouet, B., 2005. L'Acheuléen de l'ouest de la
781 France: apports du site de Menez Dregan I (Plouhinec, Finistère, France), in: Molines, N.,
782 Moncel, M.-H., Monnier, J.-L. (Eds.), *Les Premiers Peuplements en Europe*. BAR
783 International Series, Oxford, pp. 533-544.

784 Moncel, M.-H., Ashton, N., Lamotte, A., Tuffreau, A., Cliquet, D., Despriée, J., 2015. The
785 Early Acheulian of north-western Europe. *Journal of Anthropological Research* 40, 302-331.

786 Pappu, S., Gunnell, Y., Akhilesh, K., Braucher, R., Taieb, M., Demory, F., Thouveny, N.,
787 2011. Early Pleistocene presence of Acheulian hominins in south India. *Science* 331, 1596-
788 1599.

789 Parfitt, S.A., 1998. The interglacial mammalian fauna from Barnham, in: Ashton, N.M.,
790 Lewis, S.G., Parfitt, S.A. (Eds.), *Excavations at the Lower Palaeolithic Site at East Farm,*
791 *Barnham, Suffolk 1989–94*. British Museum, London, pp. 111–147.

792 Parfitt, S.A., Ashton, N.M., Lewis, S.G., Abel, R.L., Coope, R., Field, M.H., Gale, R., Hoare,
793 P.G., Larkin, N., Lewis, M.D., Karloukovski, V., Maher, B.A., Peglar, S.M., Preece, R.C.,
794 Whittaker, J.E., Stringer, C.B., 2010. Early Pleistocene Human Occupation at the Edge of the
795 Boreal Zone in Northwest Europe. *Nature* 466, 229-233.

796 Parfitt, S.A., Barendregt, R.W., Breda, M., Candy, I., Collins, M.J., Coope, G.R., Durbridge,
797 P., Field, M.H., Lee, J.R., Lister, A.M., Mutch, R., Penkman, K.E.H., Preece, R.C., Rose, J.,
798 Stringer, C.B., Symmons, R., Whittaker, J.E., Wymer, J.J., Stuart, A.J., 2005. The earliest
799 record of human activity in northern Europe. *Nature* 438, 1008–1012.

800 Parfitt, S.A., Roberts, M.B., 1999. Human modification of faunal remains, in: Roberts, M.B.,
801 Parfitt, S.A. (Eds.), *Boxgrove: A Middle Pleistocene Hominid Site at Eartham Quarry,*
802 *Boxgrove, West Sussex*. English Heritage, London, pp. 398-419.

803 Pettitt, P., White, M.J., 2012. *The British Palaeolithic: Human Societies at the Edge of the*
804 *Pleistocene World*. Routledge, London.

805 Porr, M., 2005. The making of the biface and the making of the individual, in: Gamble, C.S.,
806 Porr, M. (Eds.), *The Hominid Individual in Context: Archaeological investigations of Lower*

807 and Middle Palaeolithic landscapes, locales and artefacts Routledge, London and New York,
808 pp. 68 - 80.

809 Potts, R., 1998. Variability selection in hominid evolution. *Evolutionary Anthropology* 7, 81-
810 96.

811 Potts, R., 2013. Hominin evolution in settings of strong environmental variability. *Quaternary*
812 *Science Reviews* 73, 1-13.

813 Preece, R.C., Gowlett, J.A.J., Parfitt, S.A., Bridgland, D.R., Lewis, S.G., 2006. Humans in
814 the Hoxnian: Habitat, Context and Fire Use at Beeches Pit, West Stow, Suffolk, UK. *Journal*
815 *of Quaternary Science* 21, 485-496.

816 Rae, T.C., Koppe, T., Stringer, C.B., 2011. The Neanderthal face is not cold adapted. *Journal*
817 *of Human Evolution* 60, 234-239.

818 Ramirez Rozzi, F.V., Bermudez de Castro, J.M., 2004. Surprisingly rapid growth in
819 Neanderthals. *Nature* 428, 936-939.

820 Richter, D., Krbetschek, M., 2015. The age of the Lower Paleolithic occupation at
821 Schönningen. *Journal of Human Evolution* 89, 46-56.

822 Robson, S.L., Wood, B., 2008. Hominin life history: reconstruction and evolution. *Journal of*
823 *Anatomy* 212, 394-425.

824 Rodríguez, J., Burjachs, F., Cuenca-Bescós, G., García, N., Van der Made, J., Pérez
825 González, A., Blain, H.A., Expósito, I., López-García, J.M., García Antón, M., Allué, E.,
826 Cáceres, I., Huguet, R., Mosquera, M., Ollé, A., Rosell, J., Parés, J.M., Rodríguez, X.P.,
827 Díez, C., Rofes, J., Sala, R., Saladié, P., Vallverdú, J., Bennisar, M.L., Blasco, R., Bermúdez
828 de Castro, J.M., Carbonell, E., 2011. One million years of cultural evolution in a stable
829 environment at Atapuerca (Burgos, Spain). *Quaternary Science Reviews* 30, 1396-1412.

830 Roebroeks, W., 2001. Hominid behaviour and the earliest occupation of Europe: an
831 exploration. *Journal of Human Evolution* 41, 437-461.

832 Roebroeks, W., 2006. The human colonisation of Europe: where are we? *Journal of*
833 *Quaternary Science* 21, 425-435.

834 Roebroeks, W., van Kolfschoten, T., 1994. The earliest occupation of Europe: a short
835 chronology. *Antiquity* 68, 489-503.

836 Roebroeks, W., Villa, P., 2011. On the earliest evidence for habitual use of fire in Europe.
837 *Proceedings of the National Academy of Sciences* 108, 5209-5214.

838 Rosas, A., Ríos, L., Estalrich, A., Liversidge, H., García-Taberner, A., Huguet, R.,
839 Cardoso, H., Bastir, M., Lalueza-Fox, C., de la Rasilla, M., Dean, C., 2017. The growth
840 pattern of Neandertals, reconstructed from a juvenile skeleton from El Sidrón (Spain).
841 *Science* 357, 1282-1287.

842 Saladié, P., Huguet, R., Díez, C., Rodríguez-Hidalgo, A., Cáceres, I., Vallverdú, J., Rosell, J.,
843 Bermúdez de Castro, J.M., Carbonell, E., 2011. Carcass transport decisions in *Homo*
844 *antecessor* subsistence strategies. *Journal of Human Evolution* 61, 425-446.

845 Saragusti, I., Sharon, I., Katzenelson, O., Avnir, D., 1998. Quantitative Analysis of the
846 Symmetry of Artefacts: Lower Paleolithic Handaxes. *Journal of Archaeological Science* 25,
847 817 - 825.

848 Schwartz, G.T., 2012. Growth, Development, and Life History throughout the Evolution of
849 *Homo*. *Current Anthropology* 53, S395-S408.

850 Shettleworth, S.J., 2010. *Cognition, evolution and behavior*, 2nd ed. Oxford University Press,
851 New York.

852 Shotton, F.W., Keen, D.H., Coope, G.R., Currant, A.P., Gibbard, P.L., Aalto, M., Peglar,
853 S.M., Robinson, J.E., 1993. The Middle Pleistocene Deposits of Waverley Wood Pit,
854 Warwickshire, England. *Journal of Quaternary Science* 8, 293-325.

855 Shultz, S., Nelson, E., Dunbar, R.I.M., 2012. Hominin Cognitive Evolution: Identifying
856 Patterns and Processes in the Fossil and Archaeological Record. *Philosophical Transactions of*
857 *the Royal Society Series B* 367, 2130-2140.

858 Smith, T.M., Toussaint, M., Reid, D.J., Olejniczak, A.J., Hublin, J.-J., 2007. Rapid dental
859 development in a Middle Paleolithic Belgian Neanderthal. *Proceedings of the National*
860 *Academy of Sciences* 104, 20220-20225.

861 Stahlschmidt, M.C., Miller, C.E., Ligouis, B., Goldberg, P., Berna, F., Urban, B., Conard,
862 N.J., 2015. The depositional environments of Schöningen 13 II-4 and their archaeological
863 implications. *Journal of Human Evolution* 89, 71-91.

864 Stopp, M., 1993. Taphonomic analysis of the faunal assemblage, in: Singer, R., Gladfelter,
865 B.G., Wymer, J.J. (Eds.), *The Lower Palaeolithic Site at Hoxne, England*. University of
866 Chicago Press, Chicago, pp. 139–149.

867 Stout, D., Apel, J., Commander, J., Roberts, M., 2014. Late Acheulean technology and
868 cognition at Boxgrove, UK. *Journal of Archaeological Science* 41, 576-590.

869 Stringer, C., 2012. The Status of *Homo heidelbergensis* (Schoetensack 1908). *Evolutionary*
870 *Anthropology* 21, 101 - 107.

871 Thackeray, J.F., 2015. Estimating the age and affinities of *Homo naledi*. *South Africa Journal*
872 *of Science* 111, 1-2.

873 Toro-Moyano, I., Martínez-Navarro, B., Agustí, J., Souday, C., Bermúdez de Castro, J.M.,
874 Martínón-Torres, M., Fajardo, B., Duval, M., Falguères, C., Oms, O., Parés, J.M., Anadón,
875 P., Julià, R., García-Aguilar, J.M., Moigne, A.-M., Espigares, M.P., Ros-Montoya, S.,
876 Palmqvist, P., 2013. The oldest human fossil in Europe, from Orce (Spain). *Journal of Human*
877 *Evolution* 65, 1-9.

878 Tuffreau, A., Antoine, P., 1995. The earliest occupation of Europe: continental northwestern
879 Europe, in: Roebroeks, W., van Kolfschoten, T. (Eds.), *The Earliest Occupation of Europe*.
880 University of Leiden & European Science Foundation, Leiden, pp. 147-163.

881 Turner, A., 1992. Large Carnivores and Earliest European Hominids - Changing
882 Determinants of Resource Availability During the Lower and Middle Pleistocene. *Journal of*
883 *Human Evolution* 22, 109-126.

884 Urban, B., Bigga, G., 2015. Environmental reconstruction and biostratigraphy of late Middle
885 Pleistocene lakeshore deposits at Schöningen. *Journal of Human Evolution* 89, 57-70.

886 Van Buskirk, J., 2012. Behavioural plasticity and environmental change, in: Candolin, U.,
887 Wong, B. (Eds.), *Behavioural responses to a changing world: mechanisms and consequences*.
888 Oxford University Press, Oxford, pp. 145-158.

889 Van Kolfschoten, T., Buhrs, E., Verheijen, I., 2015a. The larger mammal fauna from the
890 Lower Paleolithic Schöningen Spear site and its contribution to hominin subsistence. *Journal*
891 *of Human Evolution* 89, 138-153.

892 Van Kolfschoten, T., Parfitt, S.A., Serangeli, J., Bello, S.M., 2015b. Lower Paleolithic bone
893 tools from the ‘Spear Horizon’ at Schöningen (Germany). *Journal of Human Evolution* 89,
894 226-263.

895 van Schaik, C.P., 2013. The costs and benefits of flexibility as an expression of behavioural
896 plasticity: a primate perspective. *Philosophical Transactions of the Royal Society B* 368.


897 Voormolen, B., 2008. Ancient hunters, modern butchers: Schöningen 13II-4, a kill-butchery
898 site dating from the northwest European Lower Palaeolithic. University of Leiden.


899 Warren, S.H., 1911. On a Palaeolithic (?) Wooden Spear. *Quarterly Journal of the Geological*
900 *Society of London* 67, xcix.

901 Wenban-Smith, F., 2004. Handaxe typology and Lower Palaeolithic cultural development:
902 flint, cleavers and two giant handaxes from Cuxton. *Lithics* 25, 11 - 21.


903 White, M.J., 1998. Twisted Ovate Bifaces in the British Lower Palaeolithic: Some
904 Observations and Implications, in: Ashton, N., Healy, F., Pettitt, P. (Eds.), *Stone Age*

- 905 Archaeology: Essays in honour of John Wymer. Lithic Studies Society Occasional Paper 6,
906 Oxbow Monograph 12, Oxford, pp. 98-104.
907 Wynn, T., 1995. Handaxe enigmas. *World Archaeology* 27, 10 - 23.
908







Winter



Spring



Summer



Autumn





Table 1: Summarising the long, short and modified short chronologies, as originally proposed (Carbonell et al., 1996; Dennell and Roebroeks, 1996; Gamble, 1999; Roebroeks and van Kolfschoten, 1994).

Model	Long Chronology (Carbonell et al., 1996)	Short Chronology (Roebroeks and van Kolfschoten, 1994)		Modified Short Chronology (Dennell and Roebroeks, 1996)
Summary	Europe occupied in the Early Pleistocene, and in some cases before c. 1.5 mya	“No undisputable proof for human occupation of Europe prior to about 500,000 years ago” (Roebroeks and van Kolfschoten, 1994: 500)		“Hominids might have occasionally moved into southern Europe well before 500,000 years ago, as and when conditions permitted” (Dennell and Roebroeks, 1996: 540)
Evidence				
Timeframe	Before 780,000 years ago (pan-Europe)	Before 500,000 years ago (pan-Europe)	After 500,000 years ago (pan-Europe)	Before 780,000 years ago (Southern Europe)
Archaeological signature	Small assemblages (typically 10s and 100s of artefacts)	Small series of isolated pieces selected from a natural pebble background	Large collections from excavated knapping floors with conjoinable material	Small assemblages (typically 10s and 100s of artefacts)
Context	Cave(?) and open-air sites (pan-Europe)	Disturbed secondary contexts	Primary context sites (fine-grained matrix)	Cave(?) and open-air sites (Iberia)

Assemblage character	Flake and pebble tools	Contested 'primitive' assemblages	Uncontested Acheulean and non-Acheulean industries	Flake and pebble tools
Human Remains	Occasional	No human remains	Human remains common	Occasional

Table 2: Key chronological phases and selected northern European sites (after Ashton, 2015).

Period	Chronology	Sites
Later Early Pleistocene	c. 1.3–0.78mya	Happisburgh III, Lunery-Rosières, Pont-de-Lavaud
Early Middle Pleistocene	c. 0.78–0.5mya	Abbeville, Boxgrove, Happisburgh I, High Lodge, Kärlich G, La Genetière, La Noira, Miesenheim, Pakefield, Warren Hill
Later Middle Pleistocene	c. 0.5–0.3mya	Beeches Pit, Bilzingsleben, Clacton, Hoxne, Schöningen, Soucy, Swanscombe,

Table 3: Prevailing habitat characteristics in Europe over the last 1.2mya (re-drawn after Kahlke et al., 2011: Fig. 6).

Region	Prevailing habitats (inferred from large mammal communities)				
Apennine Peninsula	High variety of open / forested habitats		Open woodland / steppe		
Iberian Peninsula	Open savannah / lightly forested habitats		Open woodland, tree savannah / steppe		
Western Europe	High variety of open / forested habitats		Woodland / steppe	Variety of open to forested habitats	
Northwest Europe	Woodland		Woodland with open patches / mixed habitats	Steppe / woodland	
Central Europe	High variety of open / forested habitats		Steppe / woodland with open patches	Steppe-tundra / woodland	
Eastern Europe	High variety of forest steppe / open steppe habitats		Open steppe / forest steppe	Steppe-tundra / woodland	
Chronology (mya)	1.2	1.0	0.8	0.6	0.4

Table 4: Inferred environments and key supporting evidence at Atapuerca in the Early and Middle Pleistocene (after Rodríguez et al., 2011: table 9).

Evidence	Key Patterns (Early & Middle Pleistocene)	
Small vertebrates	<ul style="list-style-type: none"> • Significant woodland component in herpetofauna • Small mammals suggest moist, open habitats 	
Large mammals	<ul style="list-style-type: none"> • All of temperate affinity • None unequivocally indicating cold or arid environments 	
Vegetation	<ul style="list-style-type: none"> • Continuous presence of Mediterranean taxa & mesic trees • One possible cold steppe landscape phase 	
Palaeotemperature (amphibian & squamate assemblages)	<ul style="list-style-type: none"> • Always warmer than Burgos today • “Cold”/“warm” period differences of only c. 2°C 	
Annual precipitation	<ul style="list-style-type: none"> • 750mm (dry)–1049mm (humid), compared to c. 570mm in present 	
Environment types	<p>Early Pleistocene</p> <ul style="list-style-type: none"> • Mediterranean open woodland • Open woodland and steppe • Open arid (?) • Open woodland and moist meadows • Temperate open woodlands and meadows with lagoons 	<p>Middle Pleistocene</p> <ul style="list-style-type: none"> • Humid temperate open woodland • Mediterranean open woodland • Moist meadows with scattered trees • Open woodland and meadows • Steppe

Table 5: Winter and summer temperature estimates for Early and Middle Pleistocene British and German sites (with selected present day equivalents for reference). ¹Sensitivity tests on coleoptera-based MCR procedures suggest that winter temperature estimates are usually too warm (Pettitt and White, 2012: 35). ²The specific source of the palaeo-temperature estimates is not stated, but the fauna includes molluscs and ostracods; ³East Anglian data based on Met Office annual mean seasonal temperatures (1910–2016; http://www.metoffice.gov.uk/pub/data/weather/uk/climate/datasets/Tmean/date/East_Anglia.txt); ⁴Bilzingsleben data based on Deutscher Wetterdienst (German Weather Service) monthly mean January and July temperatures (1951–2017; Erfurt-Weimar station; <https://www.dwd.de/DE/leistungen/klimadatendeutschland/klarchivtagmonat.html>)

Site	T _{min} (°C)	T _{max} (°C)	Evidence ¹	Age (MIS)	Source
Early Pleistocene					
Happisburgh III (Bed E)	-3 - 0	+16 - +18	Coleoptera	Late 25 or late 21	(Ashton and Lewis, 2012; Parfitt et al., 2010)
Early Middle Pleistocene					
Pakefield (Bed Cii-Ciii)	-6 - +4	+17 - +23	Coleoptera	17 or later 19	(Ashton and Lewis, 2012; Coope, 2006b)
Boxgrove (Unit 4c & Freshwater Silt Bed ≈ Units 4b & 4c)	-4 - +4	+15 - +20	Ostracods (MOTR) & Herpetofauna (MCR)	13	(Ashton and Lewis, 2012; Holman, 1999; Holmes

					et al., 2010)
Happisburgh I (Organic Mud)	-11 - -3	+12 - +15	Coleoptera	13?	(Ashton and Lewis, 2012; Coope, 2006b)
High Lodge (Bed C1)	-4 - +1	+15 - +16	Coleoptera	13?	(Coope, 2006b)
Waverley Wood (Channel 2, Organic Mud)	-	+10 - +15	Coleoptera	13 or 15	(Coope, 2006b; Shotton et al., 1993)
Brooksby (Redland's Brooksby Channel)	-10 - +2	+15 - +16	Coleoptera	13 or 15	(Coope, 2006b)
Later Middle Pleistocene					
Barnham (Unit 5c; Holl)	-	+17 - +18	Herpetofauna	11c	(Holman, 1998)
Hoxne (Stratum D; Holla)	-10 - +6	+15 - +19	Coleoptera	11c	(Ashton et al., 2008a; Coope, 1993)
Bilzingsleben II	-0.5 - +3	+20 - +25	Mollusca & ostracods ²	11	(Mania, 1995; Mania and Mania, 2003)
Schöningen 13 II-4	-4 - -1	+16	Mollusca & ostracods	9	(Urban and Bigga, 2015)

Present Day Comparisons					
East Anglia (present day) ³	-0.7 - +6.9	+14.2 - + 18.0	-	-	-
Bilzingsleben (present day) ⁴	-3.3 - +2.0	+12.4 - +22.7	-	-	-

Table 6: Palaeoenvironmental reconstructions for Early and Middle Pleistocene British sites and German sites.

Site	Palaeoenvironmental description	Age (MIS)	Source
Early Pleistocene			
Pont-de-Lavaud	Closed deciduous forest; warm, wet climate	30	(Messenger et al., 2011)
Happisburgh III (Unit C)	Cool temperate; heathland & coniferous forest	Late 25 or late 21	(Ashton and Lewis, 2012)
Happisburgh III (Units D & E)	Cool temperate; dominated by coniferous forest	Late 25 or late 21	(Ashton and Lewis, 2012)
Early Middle Pleistocene			
Pakefield (Units Aiv-Ciii)	Mediterranean-type climate; marshy ground with reed-beds & alder carr, surrounded by deciduous forest	17 or later 19	(Ashton and Lewis, 2012)
Happisburgh I (Organic Mud-Organic Silt Sand)	Cool climate; marshy edge of slow-flowing river, with surrounding coniferous forest	13?	(Ashton and Lewis, 2012)
High Lodge (Units B2-C2)	Cool-temperate climate; marshy ponds, heath & grassland on river floodplain, surrounded by pine & spruce forest	13?	(Ashton and Lewis, 2012)
Boxgrove (Unit 4b)	Coastal mudflats, with occasional drying out & spread of grassland	13	(Ashton and Lewis, 2012)
Boxgrove (Unit 4c & Q1B pond deposits)	Temperate climate, with cooler winters than present; grassland, scrub & mixed woodland, with spring-fed pools	13	(Ashton and Lewis, 2012)

Boxgrove (Unit 8)	Interstadial environment	Later 13- early 12	(Ashton and Lewis, 2012)
Boxgrove (Unit 11)	Periglacial, with ameliorations in climate	Later 13- early 12	(Ashton and Lewis, 2012)
Later Middle Pleistocene			
Bilzingsleben II	Fully temperate, with subcontinental influences; mixed oak woodland & shrub associations, alternating with steppe meadows	11	(Mania and Mania, 2003)
Barnham (Unit 5c; Holl)	Temperate climate; mix of grassland & deciduous woodland on edge of slow-flowing river	11c	(Ashton and Lewis, 2012)
Beeches Pit (Unit 4; Holl)	Fully temperate environment; spring-fed pools with deciduous woodland	11c	(Ashton and Lewis, 2012)
Beeches Pit (Unit 3b; Hollb-c)	Temperate climate; shallow to stagnant pools with calcareous grassland and surrounding open woodland	11c	(Ashton and Lewis, 2012)
Hoxne (Unit B1)	Cooler temperatures (<i>cf.</i> units D & E); probable birch & pine forest; some grassland areas	11a	(Ashton and Lewis, 2012)
Swanscombe (Lower Loam; Hollb-c)	Grassy floodplain with shallow pools & occasional flooding; mixed oak forest nearby, with hazel & alder	11c	(Ashton and Lewis, 2012)
Schöningen 13 II-4	Mosaic swamp, with reed-like and sedge vegetation; regional context: highly	9	(Stahlschmidt et al., 2015;

	continental dry steppe/open forest; tree taxa dominated by pine & birch		Urban and Bigga, 2015)
--	--	--	---------------------------

Table 7: European sites with butchery evidence.

Period	Species	Butchery Evidence	Sites
Early Pleistocene	<i>Bison</i> sp. <i>B. menneri</i> <i>B. voigtstedtensis</i> <i>C. elaphus</i> <i>C. s.l. nestii vallonnetensis</i> <i>Cervid</i> sp. <i>Cercopithecidae</i> <i>Eucladoceros</i> <i>Equus</i> sp. <i>Hippopotamus</i> <i>S. etruscus</i> <i>S. hundsheimensis</i> <i>U. dolinensis</i> <i>V. praeglacialis</i>	Defleshing; disarticulation; dismemberment; evisceration; filleting; skinning	Gran Dolina TD6 (Saladié et al., 2011) Sima del Elefante (Huguet et al., 2013)
Early Middle Pleistocene	<i>Bos</i> or <i>Bison</i> sp. <i>C. capreolus</i> <i>C. elaphus</i> <i>E. ferus</i> <i>S. hundsheimensis</i> <i>U. deningeri</i>	Cut-marks; defleshing; dismemberment; filleting; marrow bone breakage; skinning	Boxgrove (Bello et al., 2009; Parfitt and Roberts, 1999) Happisburgh I (Ashton et al., 2008b) Westbury (Andrews and Ghaleb, 1999)
Later Middle Pleistocene	<i>Bos</i> or <i>Bison</i> sp. <i>C. elaphus</i>	Boning; cut-marks; defleshing; dismemberment; filleting;	Barnham (Parfitt, 1998)

	<i>E. ferus</i>	marrow bone breakage; skinning	Schöningen (Van Kolfshoten et al., 2015a; Voormolen, 2008) Hoxne (Stopp, 1993)
--	-----------------	--------------------------------	---

Table 8: Summary table of early north-western biface sites, context, raw material, biface form and additional flake tool types (data from Moncel et al., 2015b).

Age (MIS)	Contexts	Raw Materials	Characteristic Biface Morphology	Additional Flake Tools	Sites
17-13	Diamicton; fluvial gravel; fluvial sand; fluvial silts; lagoonal silts; palaeosol	Andesite; siliceous; flint; quartzite	Pointed (incl. crude forms); ovate; biface-cleaver; cordiforms	Denticulates; notches; scrapers	Boxgrove (Q1B); Brandon Fields; Happisburgh I; High Lodge (Bed E; Sands & Gravels); La Noira; Maidscross Hill; Warren Hill; Waverley Wood
12-10	Gravel; slope deposit; fluvial sand; fluvial gravel; tufa; pool; colluvial; alluvial	Flint	Pointed (incl. crude forms); backed biface; cleaver; cordiform; ovate; sub-triangular	Denticulates; endscrapers; notches; scrapers; pebble tools; retouched flakes	Cagny-La-Garenne I (Cxb, Cxv, Lg, Lj & Ca assemblages); Cagny-La-Garenne II (13, 14 & J assemblages); Rue De Cagny (Series 3); La Celle; Barnham; Elveden; Beeches Pit; Swanscombe (LMG & UMG); Hoxne (Upper & Lower Industries); Ferme de l'Épinette (MS assemblage); St-Pierre-lès-Elbeuf